


Anti-Collagen III antibody ab7778

★★★★★ 24 Abreviews 565 References 画像数 5

製品の概要

製品名	Anti-Collagen III antibody
製品の詳細	Rabbit polyclonal to Collagen III
由来種	Rabbit
特異性	This type specific collagen antibody only recognizes 3D epitopes. Negligible cross-reactivity with Type I, II, IV, V or VI collagens. Non-specific cross reaction of anti-collagen antibodies with other human serum proteins or non-collagen extracellular matrix proteins is negligible
アプリケーション	適用あり: ICC/IF, IP, IHC-Fr, ELISA, IHC-P, ICC, WB
種交差性	交差種: Rat, Cow, Human 交差が予測される動物種: Mammals 
免疫原	Full length native protein (purified) corresponding to Collagen III aa 1-1466. Database link: P02461
ポジティブ・コントロール	Natural Cow Collagen III protein (ab7528) can be used as a positive control in WB. Human skin. Human testicle tissue.
特記事項	<u>For more protocol tips, please see: https://www.abcam.com/protocols/collagen</u> At least 11 genetically distinct gene products are collectively referred to as 'collagen types' or other proteins and proteoglycans of the extracellular matrix. In humans, collagens are composed of about 20 unique protein chains which under go various types of post-translational modifications and are ultimately assembled into a triple helix. This results in great diversity between collagen types. Collagens are highly conserved throughout evolution and are characterized by an uninterrupted "Glycine-X-Y" triplet repeat that is a necessary part of the triple helical structure. For these reasons it is often extremely difficult to generate antibodies with specificities to collagens. The development of type specific antibodies is dependent on NON-DENATURED three-dimensional epitopes. This preparation results in a native conformation of the protein. The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing. If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	Preservative: 0.01% Sodium azide Constituents: 0.8766% Sodium chloride, 0.424% Potassium phosphate
精製度	Immunogen affinity purified
特記事項 (精製)	Immunoaffinity chromatography using immobilized antigens followed by extensive cross-adsorption against other collagens, human serum proteins and non-collagen extracellular matrix proteins to remove any unwanted specificities.
ポリモノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab7778の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
ICC/IF	★★★★★ (3)	Use at an assay dependent concentration. PubMed: 18385800
IP		Use at an assay dependent concentration.
IHC-Fr	★★★★★ (5)	Use at an assay dependent concentration.
ELISA		Use at an assay dependent concentration. This product was assayed against 1.0 ug of Collagen III in a direct ELISA using Peroxidase conjugated Goat anti-Rabbit and ABTS (2,2'-azino-bis-[3-ethylbenthiiazoline-6-sulfonic acid]) as a substrate for 30 minutes at room temperature. This product can also be used with Biotin Conjugated Anti-Collagen III (ab6580) in a sandwich ELISA.
IHC-P	★★★★★ (13)	1/50 - 1/1000. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.
ICC		Use at an assay dependent concentration. PubMed: 19036760
WB	★★★★★ (3)	1/5000 - 1/10000. Predicted molecular weight: 138 kDa.
IF		Use at an assay dependent concentration.

ターゲット情報

機能	Collagen type III occurs in most soft connective tissues along with type I collagen.
関連疾患	Defects in COL3A1 are a cause of Ehlers-Danlos syndrome type 3 (EDS3) [MIM:130020]; also known as benign hypermobility syndrome. EDS is a connective tissue disorder characterized by

hyperextensible skin, atrophic cutaneous scars due to tissue fragility and joint hyperlaxity. EDS3 is a form of Ehlers-Danlos syndrome characterized by marked joint hyperextensibility without skeletal deformity.

Defects in COL3A1 are the cause of Ehlers-Danlos syndrome type 4 (EDS4) [MIM:130050]. EDS is a connective tissue disorder characterized by hyperextensible skin, atrophic cutaneous scars due to tissue fragility and joint hyperlaxity. EDS4 is the most severe form of the disease. It is characterized by the joint and dermal manifestations as in other forms of the syndrome, characteristic facial features (acrogeria) in most patients, and by proneness to spontaneous rupture of bowel and large arteries. The vascular complications may affect all anatomical areas. Defects in COL3A1 are a cause of susceptibility to aortic aneurysm abdominal (AAA) [MIM:100070]. AAA is a common multifactorial disorder characterized by permanent dilation of the abdominal aorta, usually due to degenerative changes in the aortic wall. Histologically, AAA is characterized by signs of chronic inflammation, destructive remodeling of the extracellular matrix, and depletion of vascular smooth muscle cells.

配列類似性

Belongs to the fibrillar collagen family.
Contains 1 fibrillar collagen NC1 domain.
Contains 1 VWFC domain.


翻訳後修飾

Proline residues at the third position of the tripeptide repeating unit (G-X-Y) are hydroxylated in some or all of the chains.
O-linked glycan consists of a Glc-Gal disaccharide bound to the oxygen atom of a post-translationally added hydroxyl group.

細胞内局在

Secreted > extracellular space > extracellular matrix.

画像




ab7778 staining Collagen III in human testicle tissue by Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections).

Tissue was fixed with paraformaldehyde and a heat mediated antigen retrieval step was performed using TE buffer pH 9.0. Samples were then incubated with ab7778 at a 1/200 dilution for 30 minutes at 20°C. The secondary used was an undiluted, HRP-conjugated goat anti-rabbit polyclonal.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Collagen III antibody (ab7778)

Image courtesy of Rudolf Jung by Abreview.




Western blot - Anti-Collagen III antibody (ab7778)

Anti-Collagen III antibody (ab7778) at 1/1000 dilution + Human collagen III at 0.1 μ g


Predicted band size: 138 kDa

Western Blot produced under denaturing and reducing conditions.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Collagen III antibody (ab7778)

ab7778 at 1:400 (45 min RT) showed strong staining in FFPE sections of human skin(left, dermis) with moderate to strong red staining and testis (right) where strong staining was observed within connective tissue between seminiferous tubules. The antibody showed strong extracellular staining within connective tissues across many organs with minimal background staining. Slides were steamed in 0.01 M sodium citrate buffer, pH 6.0 ([ab64214](#)) at 99-100°C - 20 minutes for antigen retrieval.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Collagen III antibody (ab7778)

Immunohistochemistry of ab7778. Tissue: right lobe of the liver section. A: Central Vein (CV) fibrosis, B: Non-fibrotic CV, C: Perisinusoidal fibrosis, D: Non-fibrotic area, E: Protoplasmic tract fibrosis, F: Septal fibrosis (arrow). Fixation: formalin fixed paraffin embedded. Antigen retrieval: not required. Primary antibody: Anti-collagen type I at 1:500 for 4°C for 24hr. Secondary antibody: Peroxidase biotin-streptavidin rabbit secondary antibody at 1:10,000 for 45 min at RT. Localization: Anti-collagen type III is intra and extracellular. Staining: 3,3'-diaminobenzidine tetrahydrochloride was used as the chromogen. Nuclei were counterstained purple with hematoxylin.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Collagen III antibody (ab7778)

Ab53088 staining Human skin ([ab30166](#)). Staining is localised to the extracellular matrix.

Left panel: with primary antibody at 1 ug/ml. Right panel: isotype control.

Sections were stained using an automated system DAKO Autostainer Plus, at room temperature. Sections were rehydrated and antigen retrieved with the DAKO 3-in-1 antigen retrieval buffer citrate pH 6.0 ([ab64214](#)) in a DAKO PT Link. Slides were peroxidase blocked in 3% H₂O₂ in methanol for 10 minutes. They were then blocked with Dako Protein block for 10 minutes (containing casein 0.25% in PBS) then incubated with primary antibody for 20 minutes and detected with Dako Envision Flex amplification kit for 30 minutes. Colorimetric detection was completed with diaminobenzidine for 5 minutes. Slides were counterstained with Haematoxylin and coverslipped under DePeX. Please note that for manual staining we recommend to optimize the primary antibody concentration and incubation time (overnight incubation), and amplification may be required.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish

- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors