

Anti-Ctip1/BCL-11A antibody [14B5] ab19487

KO 評価済

★★★★☆ 2 Abreviews 36 References 画像数 4

製品の概要

製品名	Anti-Ctip1/BCL-11A antibody [14B5]
製品の詳細	Mouse monoclonal [14B5] to Ctip1/BCL-11A
由来種	Mouse
特異性	Some optimisation may be required for detection of the target protein due to low levels of endogenous expression in some samples. Please see images below for suitable positive controls.
アプリケーション	適用あり: WB, Flow Cyt
種交差性	交差種: Mouse, Human
免疫原	Fusion protein corresponding to Ctip1/BCL-11A. Database link: Q9H165
エピトープ	Epitope is in core of CTIP1/Bcl11a (aa's 172-434).
ポジティブ・コントロール	WB: Raji whole cell lysate, Jurkat cell line, MEF cell line and HEK293T cell line. Flow Cyt: Ramos cells.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C.
バッファー	pH: 7.50 Preservative: 0.02% Sodium azide Constituent: HEPES
精製度	Multi-step Chromatography

ポリ/モノ	モノクローナル
クローン名	14B5
アイソタイプ	IgG1
軽鎖の種類	kappa

アプリケーション


The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab19487の使用に適用されます**
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
WB		1/1000. Detects a band of approximately 120 kDa (predicted molecular weight: 91 kDa).
Flow Cyt		Use 0.1µg for 10 ⁶ cells. ab170190 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.

ターゲット情報

機能	Functions as a myeloid and B-cell proto-oncogene. May play important roles in leukemogenesis and hematopoiesis. An essential factor in lymphopoiesis, is required for B-cell formation in fetal liver. May function as a modulator of the transcriptional repression activity of ARP1.
組織特異性	Expressed at high levels in brain, spleen thymus, bone marrow and testis. Expressed in CD34-positive myeloid precursor cells, B-cells, monocytes and megakaryocytes. Expression is tightly regulated during B-cell development.
関連疾患	Note=Chromosomal aberrations involving BCL11A may be a cause of lymphoid malignancies. Translocation t(2;14)(p13;q32.3) causes BCL11A deregulation and amplification.
配列類似性	Contains 6 C2H2-type zinc fingers.
翻訳後修飾	Sumoylated by SUMO1.
細胞内局在	Cytoplasm. Nucleus. Associates with the nuclear body. Colocalizes with SUMO1 and SENP2 in nuclear speckles.

画像



Lane 1: Wild-type HAP1 whole cell lysate (20 µg)


Lane 2: Empty (0 µg)

Lane 3: BCL11A (Ctip1) knockout HAP1 whole cell lysate (20 µg)

Lane 4: Raji whole cell lysate (20 µg)

Lanes 1 - 4: Merged signal (red and green). Green - ab19487 observed at 91 kDa. Red - loading control, [ab181602](#), observed at 37 kDa.

ab19487 was shown to specifically react with BCL11A (Ctip1) in HAP1 wild type cells. No band was observed when BCL11A (Ctip1) knockout cells were examined. Wild-type and BCL11A (Ctip1) knockout samples were subjected to SDS-PAGE. ab19487 and [ab181602](#) (Rabbit anti GAPDH loading control) were incubated overnight at 4°C at 1/1000 dilution and 1/10,000 dilution respectively. Blots were developed with Goat anti-Mouse IgG H&L (IRDye® 800CW) preabsorbed ([ab216772](#)) and Goat anti-Rabbit IgG H&L (IRDye® 680RD) preabsorbed ([ab216777](#)) secondary antibodies at 1/10,000 dilution for 1 hour at room temperature before imaging.



All lanes : Anti-Ctip1/BCL-11A antibody [14B5] (ab19487)

Lane 1 : C6 Cell Line

Lane 2 : MEF Cell Line

Lane 3 : HDFn Cell Line

Lane 4 : Jurkat Cell Line

Lane 5 : HEK293T Cell Line

Lysates/proteins at 20 µg per lane.


Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 91 kDa


The CTIP1 / BCL-11A protein band is indicated at 92 kDa. This antibody also identified a 55 kDa protein band in Western blotting, principally in mouse fibroblasts (MEF), but this is unrelated to.

Blocking was performed in 5% milk (in PBS). The primary antibody was diluted in 1% milk (in PBS) and incubated overnight at 4°C.



Flow Cytometry - Anti-Ctip1/BCL-11A antibody [14B5] (ab19487)

Overlay histogram showing Ramos cells stained with ab19487 (red line). The cells were fixed with 4% paraformaldehyde (10 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab19487, 0.1 µg/1x10⁶ cells) for 30 min at 22°C. The secondary antibody used was Alexa Fluor® 488 goat anti-mouse IgG (H+L) ([ab150113](#)) at 1/2000 dilution for 30 min at 22°C. Isotype control antibody (black line) was mouse IgG1 [ICIGG1] ([ab91353](#), 1 µg/1x10⁶ cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter. This antibody gave a positive signal in Ramos cells fixed with 80% methanol (5 min)/permeabilized with 0.1% PBS-Tween for 20 min used under the same conditions.



Western blot - Anti-Ctip1/BCL-11A antibody [14B5] (ab19487)

Anti-Ctip1/BCL-11A antibody [14B5] (ab19487) at 1 µg/ml + Raji (Human Burkitt's lymphoma cell line) Whole Cell Lysate at 10 µg

Secondary

Goat Anti-Mouse IgG H&L (HRP) preadsorbed ([ab97040](#)) at 1/5000 dilution

Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 91 kDa

Observed band size: 120 kDa

Additional bands at: 37 kDa, 42 kDa, 62 kDa. We are unsure as to the identity of these extra bands.

Exposure time: 20 minutes

The 120 kDa band observed is comparable to the molecular weight

seen with other commercially available antibodies to Ctip1 / BCL-11A.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors