

Anti-Cytochrome C Oxidase subunit VIc/COX6C antibody [3G5F7G3] ab110267

8 References **画像数 3**

製品の概要

製品名	Anti-Cytochrome C Oxidase subunit VIc/COX6C antibody [3G5F7G3]
製品の詳細	Mouse monoclonal [3G5F7G3] to Cytochrome C Oxidase subunit VIc/COX6C
由来種	Mouse
アプリケーション	適用あり: ICC/IF, Flow Cyt, WB
種交差性	交差種: Rat, Cow, Human
免疫原	Full length native protein (purified). This information is considered to be commercially sensitive.
ポジティブ・コントロール	Isolated mitochondria from Human heart, bovine heart, and rat heart. IF/ICC: HeLa cell line
特記事項	<p>This antibody clone is manufactured by Abcam. If you require a custom buffer formulation or conjugation for your experiments, please contact orders@abcam.com.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p> <p>Product was previously marketed under the MitoSciences sub-brand.</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C. Do Not Freeze.
バッファー	<p>pH: 7.5</p> <p>Preservative: 0.02% Sodium azide</p> <p>Constituent: HEPES buffered saline</p>
特記事項 (精製)	Near homogeneity as judged by SDS-PAGE. ab110267 was produced in vitro using hybridomas grown in serum-free medium, and then purified by biochemical fractionation.
ポリ/モノ	モノクローナル

クローン名	3G5F7G3
アイソタイプ	IgG2b
軽鎖の種類	kappa

アプリケーション


The Abpromise guarantee **Abpromise保証は、**次のテスト済みアプリケーションにおけるab110267の使用に適用されます
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ICC/IF		Use a concentration of 5 µg/ml.
Flow Cyt		Use 1µg for 10 ⁶ cells. ab170192 - Mouse monoclonal IgG2b, is suitable for use as an isotype control with this antibody.
WB		Use a concentration of 2 µg/ml. Predicted molecular weight: 9 kDa.

ターゲット情報

機能	This protein is one of the nuclear-coded polypeptide chains of cytochrome c oxidase, the terminal oxidase in mitochondrial electron transport.
配列類似性	Belongs to the cytochrome c oxidase subunit 6c family.
細胞内局在	Mitochondrion inner membrane.


画像



All lanes : Anti-Cytochrome C Oxidase subunit Vlc/COX6C antibody [3G5F7G3] (ab110267) at 2 µg/ml


- Lane 1 :** Isolated mitochondria from Human heart at 50 µg
- Lane 2 :** Isolated mitochondria from Bovine heart at 1 µg
- Lane 3 :** Isolated mitochondria from Rat heart at 10 µg

Predicted band size: 9 kDa



Immunocytochemistry/ Immunofluorescence - Anti-Cytochrome C Oxidase subunit Vlc/COX6C antibody [3G5F7G3] (ab110267)

ICC/IF image of ab110267 stained HeLa cells. The cells were 4% formaldehyde fixed (10 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab110267, 5µg/ml) overnight at +4°C. The secondary antibody (green) was **ab96879**, DyLight® 488 goat anti-mouse IgG (H+L) used at a 1/250 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.



Flow Cytometry - Anti-Cytochrome C Oxidase subunit Vlc/COX6C antibody [3G5F7G3] (ab110267)

Overlay histogram showing HepG2 cells stained with ab110267 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab110267, 1µg/1x10⁶ cells) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-mouse IgG (H+L) (**ab96879**) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was mouse IgG2b [PLPV219] (**ab91366**, 2µg/1x10⁶ cells) used under the same conditions. Acquisition of >5,000 events was performed. This antibody gave a positive signal in HepG2 cells fixed with 4% paraformaldehyde (10 min)/permeabilized with 0.1% PBS-Tween for 20 min used under the same conditions.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors