


Anti-Cytokeratin 13 antibody [AE8] ab16112

KO 評価済

★★★★★ **4 Abreviews** **23 References** **画像数 6**

製品の概要

製品名	Anti-Cytokeratin 13 antibody [AE8]
製品の詳細	Mouse monoclonal [AE8] to Cytokeratin 13
由来種	Mouse
アプリケーション	適用あり: IHC-Fr, ICC/IF, WB, IHC-P
種交差性	交差種: Human 交差が予測される動物種: Mouse, Rabbit 
免疫原	Full length native protein (purified). This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: A431 whole cell lysate. ICC/IF: A431 cells. IHC-P: Human tonsil. IHC-Fr: Human tonsil.
特記事項	<p>This antibody clone is manufactured by Abcam. If you require a custom buffer formulation or conjugation for your experiments, please contact orders@abcam.com.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.
バッファー	Preservative: 0.02% Sodium azide Constituents: PBS, 6.97% L-Arginine
精製度	Protein G purified
一次抗体 備考	This antibody is specific for Cytokeratin 13, which is a marker for oesophageal type differentiation which is expressed by various internal stratified epithelia.

ポリモノ	モノクローナル
クローン名	AE8
ミエローマ	P3-X63 Ag8.3
アイソタイプ	IgG
軽鎖の種類	kappa

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab16112の使用に適用されず**
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-Fr		Use a concentration of 1 µg/ml.
ICC/IF	★★★★★ (1)	Use a concentration of 1 - 5 µg/ml. PubMed: 25076852
WB		Use a concentration of 1 µg/ml. Predicted molecular weight: 50 kDa.
IHC-P	★★★★★ (3)	Use a concentration of 0.05 µg/ml.

ターゲット情報

組織特異性

Expressed in some epidermal sweat gland ducts (at protein level) and in exocervix, esophagus and placenta.

関連疾患

Defects in KRT13 are a cause of white sponge nevus of cannon (WSN) [MIM:193900]. WSN is a rare autosomal dominant disorder which predominantly affects non-cornified stratified squamous epithelia. Clinically, it is characterized by the presence of soft, white, and spongy plaques in the oral mucosa. The characteristic histopathologic features are epithelial thickening, parakeratosis, and vacuolization of the suprabasal layer of oral epithelial keratinocytes. Less frequently the mucous membranes of the nose, esophagus, genitalia and rectum are involved.


配列類似性

Belongs to the intermediate filament family.

翻訳後修飾

O-glycosylated; glycans consist of single N-acetylglucosamine residues.

画像



Western blot - Anti-Cytokeratin 13 antibody [AE8] (ab16112)

All lanes : Anti-Cytokeratin 13 antibody [AE8] (ab16112) at 1 µg/ml

Lane 1 : Wild-type A431 cell lysate

Lane 2 : KRT13 knockout A431 cell lysate


Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.

Predicted band size: 50 kDa

Observed band size: 51 kDa

False colour image of Western blot: Anti-Cytokeratin 13 antibody [AE8] staining at 1 µg/ml, shown in green; Rabbit Anti-GAPDH antibody [EPR16891] ([ab181602](#)) loading control staining at 1/20000 dilution, shown in red. In Western blot, ab16112 was shown to bind specifically to Cytokeratin 13. A band was observed at 51 kDa in wild-type A431 cell lysates with no signal observed at this size in Krt13 knockout cell line [ab269483](#) (knockout cell lysate [ab269647](#)). To generate this image, wild-type and Krt13 knockout A431 cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3 % milk in TBS-0.1 % Tween[®] 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Mouse IgG H&L (IRDye[®] 800CW) preabsorbed ([ab216772](#)) and Goat anti-Rabbit IgG H&L (IRDye[®] 680RD) preabsorbed ([ab216777](#)) at 1/20000 dilution.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 13 antibody [AE8] (ab16112)

IHC image of Cytokeratin 13 staining in a section of formalin-fixed paraffin-embedded normal human tonsil* performed on a Leica BOND™ system using the standard protocol F.

The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab16112, 0.05 µg/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with hematoxylin and mounted with DPX. The inset secondary-only control image is taken from an identical assay without primary antibody.


For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.

**Tissue obtained from the Human Research Tissue Bank, supported by the NIHR Cambridge Biomedical Research Centre*



Immunohistochemistry (Frozen sections) - Anti-Cytokeratin 13 antibody [AE8] (ab16112)

IHC image of Cytokeratin 13 staining in a section of frozen normal human tonsil*. The section was fixed using 10% formaldehyde in 1XPBS for 10 minutes. No antigen retrieval step was performed prior to staining. Non-specific protein-protein interactions were then blocked in TBS containing 0.025% (v/v) Triton X-100, 0.3M glycine and 1% (w/v) BSA for 1h at room temperature. The section was then incubated overnight at +4°C in TBS containing 0.025% (v/v) Triton X-100 and 1% (w/v) BSA with ab16112 at 1µg/ml. The section was then incubated with **ab150117** (Goat Anti-Mouse IgG H&L (Alexa Fluor® 488), 1/1000)) (shown in green) for 1 hour at room temperature. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM. The secondary-only control insert image is taken from an identical assay without primary antibody. The section was then mounted using Fluoromount®. Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8). For IHC staining systems (automated and non-automated), customers should optimize variable parameters such as antibody concentrations and incubation times. *Tissue obtained from the Human Research Tissue Bank, supported by the NIHR Cambridge Biomedical Research Centre.




Western blot - Anti-Cytokeratin 13 antibody [AE8] (ab16112)

Anti-Cytokeratin 13 antibody [AE8] (ab16112) at 1 µg/ml + A431 whole cell lysate at 20 µg

Performed under reducing conditions.

Predicted band size: 50 kDa


This blot was produced using a 4-12% Bis-tris under the MOPS buffer system. The gel was run at 200V for 55 minutes before being transferred onto a Nitrocellulose membrane at 30V for 70 minutes. The membrane was blocked for an hour using 3% milk before ab16112 and **ab181602** (Rabbit anti-GAPDH loading control) were incubated overnight at 4°C at a 1µg/ml concentration and 1/20000 dilution respectively. Antibody binding was detected using Goat anti-Mouse IgG H&L (IRDye® 800CW) preadsorbed (**ab216772**) and Goat anti-Rabbit IgG H&L (IRDye® 680RD) preadsorbed (**ab216777**) secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.



Immunocytochemistry/ Immunofluorescence - Anti-Cytokeratin 13 antibody [AE8] (ab16112)

ab16112 staining Cytokeratin 13 in A431 cells. The cells were fixed with Methanol (5min), permeabilized with 0.1%PBS-Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1%PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab16112 at 5µg/ml and **ab6046**, Rabbit polyclonal to beta Tubulin - Loading Control, at 1/1000 dilution. Cells were then incubated with **ab150117**, Goat Anti-Mouse IgG H&L (Alexa Fluor® 488) at 1/1000 dilution (shown in green) and **ab150080**, Goat polyclonal Secondary Antibody to Rabbit IgG - H&L (Alexa Fluor® 594) at 1/1000 dilution (shown in pseudocolour red). Nuclear DNA was labelled with DAPI (shown in blue).

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 13 antibody [AE8] (ab16112)

This image is courtesy of an anonymous Abreview

ab16112 staining Cytokeratin 13 in Human pharynx tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and antigen retrieval was by heat mediation in Tris pH9. Samples were incubated with primary antibody (undiluted) for 1 hour at 20°C. An undiluted HRP-conjugated Goat anti-mouse IgG polyclonal was used as the secondary antibody.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors