


Anti-DAPK2 antibody [EP1633Y] ab51601

RabMAb

★★★★★ [1 Abreviews](#) [4 References](#) [画像数 3](#)

製品の概要

製品名	Anti-DAPK2 antibody [EP1633Y]
製品の詳細	Rabbit monoclonal [EP1633Y] to DAPK2
由来種	Rabbit
アプリケーション	適用あり: Flow Cyt (Intra), WB, IHC-P 適用なし: ICC/IF or IP
種交差性	交差種: Rat, Human 交差が予測される動物種: Mouse 
免疫原	Synthetic peptide within Human DAPK2 aa 1-100 (N terminal). The exact sequence is proprietary. Database link: Q9UIK4
ポジティブ・コントロール	WB: Rat spleen lysate IHC-P: Human liver carcinoma. Flow Cyt (intra): HeLa cells.
特記事項	Our RabMAb [®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents . The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing. If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.
バッファー	pH: 7.20 Preservative: 0.05% Sodium azide Constituents: 0.1% BSA, 40% Glycerol (glycerin, glycerine), 9.85% Tris glycine, 50% Tissue culture supernatant
精製度	Tissue culture supernatant

ポリ/モノ	モノクローナル
クローン名	EP1633Y
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab51601の使用に適用されます**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。


アプリケーション	Abreviews	特記事項
Flow Cyt (Intra)		1/1000. ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
WB	★★★★★ (1)	1/1000 - 1/10000. Detects a band of approximately 42 kDa (predicted molecular weight: 42 kDa).
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

追加情報 Is unsuitable for ICC/IF or IP.

ターゲット情報

機能	Calcium/calmodulin-dependent serine/threonine kinase which acts as a positive regulator of apoptosis.
組織特異性	Ubiquitously expressed in all tissue types examined. High levels in heart, lung and skeletal muscle.
配列類似性	Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family. DAP kinase subfamily. Contains 1 protein kinase domain.
細胞内局在	Cytoplasm.

画像



Western blot - Anti-DAPK2 antibody [EP1633Y] (ab51601)


Anti-DAPK2 antibody [EP1633Y] (ab51601) at 1/2000 dilution +
Rat spleen lysate at 10 µg

Secondary

Goat anti-rabbit HRP at 1/2000 dilution


Predicted band size: 42 kDa

Observed band size: 42 kDa



Flow Cytometry (Intracellular) - Anti-DAPK2 antibody [EP1633Y] (ab51601)

Overlay histogram showing HeLa cells stained with ab51601 (red line). The cells were fixed with 4% paraformaldehyde (10 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab51601, 1/1000 dilution) for 30 min at 22°C. The secondary antibody used was Alexa Fluor® 488 goat anti-rabbit IgG (H+L) ([ab150077](#)) at 1/2000 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (0.1 µg/1x10⁶ cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter. This antibody gave a positive signal in HeLa cells fixed with 80% methanol (5 min)/permeabilized with 0.1% PBS-Tween for 20 min used under the same conditions.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-DAPK2 antibody [EP1633Y] (ab51601)

Immunohistochemical staining of paraffin embedded human liver carcinoma using ab51601 (1/50). Secondary stained using Goat anti-rabbit HRP (1/2000).

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors