# abcam

### Product datasheet

## Anti-Desmoglein 3/PVA antibody [5G11] ab14416

★★★★★ 2 Abreviews 5 References 画像数 2

#### 製品の概要

製品名 Anti-Desmoglein 3/PVA antibody [5G11]

製品の詳細 Mouse monoclonal [5G11] to Desmoglein 3/PVA

由来種 Mouse

 アプリケーション
 適用あり: IHC-P

 種交差性
 交差種: Human

免疫原 Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.

ポジティブ・コントロール IHC-P: FFPE human skin melanoma tissue sections.

特記事項 The clone number has been updated from 3G133 to 5G11 since both name the same clone and

5G11 is the original clone number.

For maximum recovery of product, centrifuge the original vial after thawing and prior to removing

the cap. Further dilutions can be made in assay buffer.

This antibody clone is manufactured by Abcam. If you require a custom buffer formulation or

conjugation for your experiments, please contact orders@abcam.com.

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

#### 製品の特性

**製品の状態** Liquid

保存方法 Shipped at 4°C. Add glycerol to a final volume of 50% for extra stability and aliquot. Store at -

20°C. Avoid freeze / thaw cycle.

**バッファー** pH: 7.40

Preservative: 0.02% Sodium azide Constituents: PBS, 6.97% L-Arginine

精製度 Protein G purified

1

**ポリ**Æ**ノ** モノクローナル **クローン名** 5G11

アイソタイプ IgG1

#### アプリケーション

**The Abpromise guarantee** <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab14416の使用に適用されますアプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-P	<b>★★★★★ (1)</b>	Use a concentration of 5 µg/ml. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

#### ターゲット情報

機能 Component of intercellular desmosome junctions. Involved in the interaction of plaque proteins

and intermediate filaments mediating cell-cell adhesion.

組織特異性 Epidermis, tongue, tonsil, esophagus and carcinomas.


**配列類似性** Contains 4 cadherin domains.

ドメイン Three calcium ions are usually bound at the interface of each cadherin domain and rigidify the

connections, imparting a strong curvature to the full-length ectodomain.

**細胞内局在** Cell membrane. Cell junction > desmosome.


#### 画像



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Desmoglein 3/PVA antibody [5G11] (ab14416)

IHC image of Desmoglein 3/PVA staining in Human skin melanoma formalin fixed paraffin embedded tissue section, performed on a Leica Bond™ system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab14416, 5µg/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Desmoglein 3/PVA antibody [5G11] (ab14416)

This image is courtesy of an anonymous Abreview

Immunohistochemical analysis of Human tongue tissue, staining Desmoglein 3/PVA with ab14416.

Tissue was fixed with formaldehyde and antigen retrieval was by proteinase K. Samples were incubated with primary antibody (1/500 in Tris buffered saline with 0.1% Tween 20) for 18 hours at 4°C. An HRP-conjugated goat anti-mouse polyclonal lgG (1/2) was used as the secondary antibody.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.co.jp/abpromise">https://www.abcam.co.jp/abpromise</a> or contact our technical team.

#### Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors