

# Anti-DMPO Nitron Adduct antibody [N1664A] ab104902

3 References [画像数 2](#)

### 製品の概要

製品名	Anti-DMPO Nitron Adduct antibody [N1664A]
製品の詳細	Mouse monoclonal [N1664A] to DMPO Nitron Adduct
由来種	Mouse
特異性	Detects ~90kDa. Recognizes DMPO, DMPO-octanoic acid, DMPO-protein adducts and DMPO-DNA adducts. Does not cross-react with non-adducted proteins or DNA.
アプリケーション	<b>適用あり:</b> WB, ELISA, ICC/IF, IP
種交差性	<b>交差種:</b> Species independent
免疫原	5,5-dimethyl-2-(8-octanoic acid)-1-pyrrolone-N-oxide conjugated to Ovalbumin.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
バッファー	Preservative: 0.05% Sodium azide Constituents: 50% Glycerol (glycerin, glycerine), Tris buffered saline
精製度	Protein G purified
ポリ/モノ	モノクローナル
クローン名	N1664A
アイソタイプ	IgG1

### アプリケーション

The Abpromise guarantee

Abpromise保証は、次のテスト済みアプリケーションにおけるab104902の使用に適用されます


アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
WB		Use a concentration of 1 - 10 µg/ml.
ELISA		Use a concentration of 1 - 10 µg/ml.
ICC/IF		Use a concentration of 1 - 10 µg/ml.
IP		Use at an assay dependent dilution. Follow the method described by Chatterjee, S. <i>et. al.</i> PubMed 19049863.


## ターゲット情報

**関連性** For measurement of protein radicals; some proteins with endogenous peroxidase activity are especially susceptible to forming DMPO adducts.

## 画像



Immunocytochemistry/Immunofluorescence analysis of Mouse macrophage cells labelling DMPO Nitron Adduct with ab104902 at 10µg/mL. Left - with DMPO, Right - no DMPO. Top to bottom - DMPO, DAPI, overlay.



**All lanes :** Anti-DMPO Nitron Adduct antibody [N1664A] (ab104902) at 10 µg/ml

**Lane 1 :** 10µM Hb

**Lane 2 :** 10µM Hb + 100µM HOCL

**Lane 3 :** 10µM Hb + 500µM HOCL

**Lane 4 :** 10µM Hb + 1000µM HOCL

**Lane 5 :** 10µM Hb with 20mM DMPO

**Lane 6 :** 10µM Hb + 100µM HOCL with 20mM DMPO

**Lane 7 :** 10µM Hb + 500µM HOCL with 20mM DMPO

**Lane 8 :** 10µM Hb + 1000µM HOCL with 20mM DMPO

**Lane 9 :** 10µM Hb with 100mM DMPO

**Lane 10 :** 10µM Hb + 100µM HOCL with 100mM DMPO

**Lane 11 :** 10µM Hb + 500µM HOCL with 100mM DMPO

**Lane 12 :** 10µM Hb + 1000µM HOCL with 100mM DMPO

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors