abcam

Product datasheet

Anti-DPP8 antibody - Aminoterminal end ab42075

* ★ ★ ★ ★ ↑ 2 Abreviews 3 References

製品の概要

製品名 Anti-DPP8 antibody - Aminoterminal end

製品の詳細 Rabbit polyclonal to DPP8 - Aminoterminal end

由来種 Rabbit

特異性 This antibody recognizes all four forms of DPP8. This antibody does not recognize other DPP

family members.

適用あり: WB アプリケーション 交差種: Human 種交差性

交差が予測される動物種: Mouse, Rat 4

免疫原 Synthetic peptide corresponding to the aminoterminal end of human DPP8. Immunogen の所有権

に関して (Peptide available as ab44587.)

特記事項

This antibody recognizes all four forms of DPP8

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

バッファー Preservative: 0.05% Sodium azide

Constituents: PBS, 50% Glycerol, 2.9% Sodium chloride

精製度 Immunogen affinity purified

一次抗体 備考 This antibody recognizes all four forms of DPP8

ポリクローナル ポリモノ

アイソタイプ ΙgG

アプリケーション

The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab42075の使用に適用されますアプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB	★★★★☆ (2)	Use at an assay dependent concentration. Predicted molecular weight: 103 kDa. Use at 1/1000 when using colourimetric substrates, use at 1/5000 when using chemiluminescent substrates. Dilution optimised using Chromogenic detection.

ターゲット情報

機能	Dipeptidyl peptidase that cleaves off N-terminal dipeptides from proteins having a Pro or Ala residue at position 2. May play a role in T-cell activation and immune function.	
組織特異性	Ubiquitously expressed, with highest levels in testis, placenta, prostate, muscle and brain.	
配列類似性	Belongs to the peptidase S9B family. DPPIV subfamily.	
細胞内局在	Cytoplasm.	

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors