


# Anti-DRIP130 antibody ab70449

[1 References](#) [画像数 1](#)

### 製品の概要

---

製品名	Anti-DRIP130 antibody
製品の詳細	Rabbit polyclonal to DRIP130
由来種	Rabbit
アプリケーション	<b>適用あり:</b> IP
種交差性	<b>交差種:</b> Human <b>交差が予測される動物種:</b> Mouse, Horse, Guinea pig, Dog, Chimpanzee, Ferret, Rhesus monkey, Gorilla, Orangutan, Zebra finch 
免疫原	Synthetic peptide, corresponding to a region between amino acids 400-450 of Human DRIP130 (SwissProt Q9ULK4)
ポジティブ・コントロール	Nuclear extract from HeLa cells
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

### 製品の特性

---

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
バッファー	pH: 7 Preservative: 0.09% Sodium azide Constituents: 1.815% Tris, 1.764% Sodium citrate, 0.021% PBS
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

**The Abpromise guarantee**      **Abpromise保証は、次のテスト済みアプリケーションにおけるab70449の使用に適用されます**  
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
IP		

**追加情報**

IP: Use at 1 - 4 µg/mg of lysate.

Is unsuitable for WB.

Not yet tested in other applications.

Optimal dilutions/concentrations should be determined by the end user.

**ターゲット情報**

**機能**

Required for transcriptional activation subsequent to the assembly of the preinitiation complex (By similarity). Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors. Required for transcriptional activation by adenovirus E1A protein. Required for ELK1-dependent transcriptional activation in response to activated Ras signaling.


**配列類似性**

Belongs to the Mediator complex subunit 23 family.

**細胞内局在**

Nucleus.

**画像**



Detection of Human DRIP130 by Immunoprecipitation in Nuclear extract (5 mg) from HeLa cells using ab70449 at 4 µg/5 mg extract.  
 Detection: Coomassie stain.

Immunoprecipitation - Anti-DRIP130 antibody  
 (ab70449)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

## **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors