

Anti-DUOX2 antibody ab97266

5 References [画像数 2](#)

製品の概要

製品名	Anti-DUOX2 antibody
製品の詳細	Rabbit polyclonal to DUOX2
由来種	Rabbit
特異性	From Jan 2024, QC testing of replenishment batches of this polyclonal changed. All tested and expected application and reactive species combinations are still covered by our Abcam product promise. However, we no longer test all applications. For more information on a specific batch, please contact our Scientific Support who will be happy to help.
アプリケーション	適用あり: WB, IHC-P
種交差性	交差種: Rat 交差が予測される動物種: Human, Chimpanzee, Macaque monkey, Orangutan 
免疫原	Synthetic peptide corresponding to Rat DUOX2 aa 1-100 conjugated to keyhole limpet haemocyanin. Database link: Q9ES45
ポジティブ・コントロール	This antibody gave a positive signal in Rat Colon tissue lysate. This antibody gave a positive result in IHC in the following FFPE tissue: Rat Thyroid.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	pH: 7.40 Preservative: 0.02% Sodium azide Constituent: PBS

Batches of this product that have a concentration < 1mg/ml may have BSA added as a stabilising agent. If you would like information about the formulation of a specific lot, please contact our scientific support team who will be happy to help.

精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション


The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab97266の使用に適用されます**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		Use a concentration of 1 µg/ml. Predicted molecular weight: 175 kDa.
IHC-P		Use a concentration of 1 µg/ml. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

ターゲット情報

機能	Generates hydrogen peroxide which is required for the activity of thyroid peroxidase/TPO and lactoperoxidase/LPO. Plays a role in thyroid hormones synthesis and lactoperoxidase-mediated antimicrobial defense at the surface of mucosa. May have its own peroxidase activity through its N-terminal peroxidase-like domain.
組織特異性	Expressed in colon, small intestine, duodenum and tracheal surface epithelial cells (at protein level). Expressed in thyrocytes. Also detected in kidney, liver, lung, pancreas, prostate, salivary glands, rectum and testis.
パスウェイ	Hormone biosynthesis; thyroid hormone biosynthesis.
関連疾患	Defects in DUOX2 are a cause of thyroid dysmorphogenesis 6 (TDH6) [MIM:607200]. A disorder due to a defective conversion of accumulated iodide to organically bound iodine. The iodide organification defect can be partial or complete.
配列類似性	In the N-terminal section; belongs to the peroxidase family. Contains 3 EF-hand domains. Contains 1 FAD-binding FR-type domain. Contains 1 ferric oxidoreductase domain.
発生段階	Widely expressed in fetal tissues.
翻訳後修飾	N-glycosylated.
細胞内局在	Apical cell membrane. Localizes to the apical membrane of epithelial cells.

画像



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-DUOX2 antibody (ab97266)

IHC image of DUOX2 staining in Rat thyroid formalin fixed paraffin embedded tissue section, performed on a Leica Bond™ system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab97266, 1 µg/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.



Western blot - Anti-DUOX2 antibody (ab97266)

Anti-DUOX2 antibody (ab97266) at 1 µg/ml (Milk blocking (3%)) + Colon (Rat) Tissue Lysate at 25 µg

Secondary

Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/10000 dilution

Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 175 kDa

Observed band size: 171 kDa

Exposure time: 20 minutes

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise,

please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors