abcam

Product datasheet

DyLight® 488 Anti-HA tag antibody [16B12] ab117488

★★★★★ 1 Abreviews 1 References 画像数 1

製品の概要

製品名 DyLight® 488 Anti-HA tag antibody [16B12]

製品の詳細 DyLight® 488 Mouse monoclonal [16B12] to HA tag

由来種 Mouse

標識 DyLight® 488. Ex: 493nm, Em: 518nm

アプリケーション 適用あり: ICC/IF, Flow Cyt

種交差性 交差種: Species independent

免疫原 Synthetic peptide corresponding to HA tag.

特記事項 The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C.

バッファー pH: 7.40

Preservative: 0.02% Sodium azide

Constituents: 1.6% Sodium phosphate, 0.88% Sodium chloride, 1% BSA

精製度 Protein G purified

ポリ/モノ モノクローナル

クローン名 16B12 アイソタイプ laG1

アプリケーション

アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。


アプリケーション	Abreviews	特記事項
ICC/IF	★★★★ <u>(1)</u>	1/500 - 1/5000.
Flow Cyt		Use at an assay dependent concentration.

ターゲット情報

関連性

Human influenza hemagglutinin (HA) is a surface glycoprotein required for the infectivity of the human virus. The HA tag is derived from the HA molecule corresponding to amino acids 98-106 has been extensively used as a general epitope tag in expression vectors. Many recombinant proteins have been engineered to express the HA tag, which does not appear to interfere with the bioactivity or the biodistribution of the recombinant protein. This tag facilitates the detection, isolation, and purification of the proteins.

画像



ab117488 staining HA tag in Monkey cos7 cells by ICC/IF (Immunocytochemistry/immunofluorescence). Cells were fixed with formaldehyde. Samples were incubated with primary antibody (1/100) for 1 hour at 22°C.

Immunocytochemistry/ Immunofluorescence -DyLight® 488 Anti-HA tag antibody [16B12] (ab117488)

This image is courtesy of an Abreview submitted by Ya Zhou

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors