

# Anti-ECSIT antibody ab21288

**7 References**   [画像数 1](#)

### 製品の概要

製品名	Anti-ECSIT antibody
製品の詳細	Rabbit polyclonal to ECSIT
由来種	Rabbit
アプリケーション	<b>適用あり:</b> IHC-P, IP, ICC/IF, WB
種交差性	<b>交差種:</b> Mouse, Human
免疫原	Synthetic peptide corresponding to 14 amino acids near the C-terminus of ECSIT (Human).
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C.
バッファー	pH: 7.2 Preservative: 0.02% Sodium azide Constituent: PBS
精製度	Affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

### アプリケーション


**The Abpromise guarantee**   **Abpromise保証は、** 次のテスト済みアプリケーションにおけるab21288の使用に適用されます  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-P		Use at an assay dependent concentration.
IP		Use at an assay dependent concentration. PubMed: 21163940
ICC/IF		Use at an assay dependent concentration. PubMed: 21163940
WB		Use a concentration of 0.5 - 1 µg/ml. Detects a band of approximately 42 kDa (predicted molecular weight: 49 kDa).

## ターゲット情報

機能	Adapter protein of the Toll-like and IL-1 receptor signaling pathway that is involved in the activation of NF-kappa-B via MAP3K1. Promotes proteolytic activation of MAP3K1. Involved in the BMP signaling pathway. Required for normal embryonic development (By similarity). Required for efficient assembly of mitochondrial NADH:ubiquinone oxidoreductase.
配列類似性	Belongs to the ECSIT family.
細胞内局在	Cytoplasm. Nucleus. Mitochondrion.

## 画像



**Lane 1 :** Anti-ECSIT antibody (ab21288) at 0.5 µg/ml

**Lane 2 :** Anti-ECSIT antibody (ab21288) at 1 µg/ml

**Lane 3 :** Anti-ECSIT antibody (ab21288) at 2 µg/ml

**All lanes :** human heart lysate

**Predicted band size:** 49 kDa

**Observed band size:** 42 kDa

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

#### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors