abcam

Product datasheet

Anti-EGFR (phospho Y1086) antibody [EP2258IHCY] ab68470

ועלשעבע RabMAb

画像数3

製品の概要

製品名 Anti-EGFR (phospho Y1086) antibody [EP2258IHCY]

製品の詳細 Rabbit monoclonal [EP2258IHCY] to EGFR (phospho Y1086)

由来種 Rabbit

特異性 Recognises EGFR phosphorylated on Tyrosine 1086 of the mature human isoform 1

(corresponding to Y1110 from the precursor form P00533-1/p170)

アプリケーション 適用あり: ICC/IF, WB

適用なし: Flow Cyt or IHC-P

種交差性 交差種: Human

免疫原 Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

ポジティブ・コントロール A431 cell lysate, A431 cells.

特記事項 This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity

- Long-term security of supply

- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with

these species. Please contact us for more information.

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

バッファー pH: 7.2

Preservative: 0.01% Sodium azide

Constituents: 9% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA, 50% Tissue culture

supernatant

1

精製度 Protein A purified

ポリ/モノ モノクローナル

クローン名 EP2258IHCY

アイソタイプ IgG

アプリケーション

The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab68470の使用に適用されます

アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ICC/IF		
WB		

追加情報 ICC/IF: 1/100 - 1/250.

IP: 1/40.

WB: 1/1000 - 1/5000. Detects a band of approximately 200 kDa (predicted molecular weight: 134

kDa).

Is unsuitable for Flow Cyt or IHC-P.

Not yet tested in other applications.

Optimal dilutions/concentrations should be determined by the end user.

ターゲット情報

機能 Receptor tyrosine kinase binding ligands of the EGF family and activating several signaling

cascades to convert extracellular cues into appropriate cellular responses. Known ligands include EGF, TGFA/TGF-alpha, amphiregulin, epigen/EPGN, BTC/betacellulin, epiregulin/EREG and HBEGF/heparin-binding EGF. Ligand binding triggers receptor homo- and/or heterodimerization and autophosphorylation on key cytoplasmic residues. The phosphorylated receptor recruits adapter proteins like GRB2 which in turn activates complex downstream signaling cascades. Activates at least 4 major downstream signaling cascades including the RAS-RAF-MEK-ERK, PI3 kinase-AKT, PLCgamma-PKC and STATs modules. May also activate the NF-kappa-B signaling cascade. Also directly phosphorylates other proteins like RGS16, activating its GTPase activity and probably coupling the EGF receptor signaling to the G protein-coupled receptor signaling. Also phosphorylates MUC1 and increases its interaction with SRC and CTNNB1/beta-

catenin.

Isoform 2 may act as an antagonist of EGF action.

組織特異性 Ubiquitously expressed. Isoform 2 is also expressed in ovarian cancers.

関連疾患 Lung cancer

Inflammatory skin and bowel disease, neonatal, 2

配列類似性 Belongs to the protein kinase superfamily. Tyr protein kinase family. EGF receptor subfamily.

Contains 1 protein kinase domain.

翻訳後修飾 Phosphorylation at Ser-695 is partial and occurs only if Thr-693 is phosphorylated.

Phosphorylation at Thr-678 and Thr-693 by PRKD1 inhibits EGF-induced MAPK8/JNK1 activation. Dephosphorylation by PTPRJ prevents endocytosis and stabilizes the receptor at the plasma membrane. Autophosphorylation at Tyr-1197 is stimulated by methylation at Arg-1199 and enhances interaction with PTPN6. Autophosphorylation at Tyr-1092 and/or Tyr-1110 recruits STAT3. Dephosphorylated by PTPN1 and PTPN2.

Monoubiquitinated and polyubiquitinated upon EGF stimulation; which does not affect tyrosine kinase activity or signaling capacity but may play a role in lysosomal targeting. Polyubiquitin linkage is mainly through 'Lys-63', but linkage through 'Lys-48', 'Lys-11' and 'Lys-29' also occurs. Deubiquitination by OTUD7B prevents degradation. Ubiquitinated by RNF115 and RNF126. Methylated. Methylation at Arg-1199 by PRMT5 stimulates phosphorylation at Tyr-1197.

Secreted and Cell membrane. Endoplasmic reticulum membrane. Golgi apparatus membrane. Nucleus membrane. Endosome membrane. Nucleus. In response to EGF, translocated from the cell membrane to the nucleus via Golgi and ER. Endocytosed upon activation by ligand. Colocalized with GPER1 in the nucleus of estrogen agonist-induced cancer-

associated fibroblasts (CAF).

細胞内局在

画像

1 2
25015010075503725201510
Beta Tubulin

Western blot - Anti-EGFR (phospho Y1086) antibody [EP2258IHCY] (ab68470) **All lanes :** Anti-EGFR (phospho Y1086) antibody [EP2258IHCY] (ab68470) at 1/5000 dilution

Lane 1: A431 cell lysate, untreated

Lane 2: A431 cell lysate, treated with EGF

Lysates/proteins at 10 µg per lane.


Secondary

All lanes: HRP conjugated Goat anti-rabbit at 1/2000 dilution

Predicted band size: 134 kDa

Observed band size: 200 kDa

Immunocytochemistry/ Immunofluorescence - Anti-EGFR (phospho Y1086) antibody [EP2258IHCY] (ab68470) A431 cells stained with ab68470 at 1/100 dilution.



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors