




# Anti-Eph receptor B1/NET (phospho Y317) antibody ab30562

### 製品の概要

製品名	Anti-Eph receptor B1/NET (phospho Y317) antibody
製品の詳細	Rabbit polyclonal to Eph receptor B1/NET (phospho Y317)
由来種	Rabbit
特異性	ab30562 recognises EphB1 (phospho Y317).
アプリケーション	<b>適用あり:</b> WB
種交差性	<b>交差種:</b> Rat <b>交差が予測される動物種:</b> Mouse, Chicken, Cow, Human, Xenopus laevis, Zebrafish 
免疫原	Synthetic peptide corresponding to Chicken Eph receptor B1/NET (phospho Y317).  <a href="#">Run BLAST with ExPASy</a>  <a href="#">Run BLAST with NCBI</a>
ポジティブ・コントロール	Rat testes lysate
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
バッファー	pH: 7.50 Constituents: 0.238% HEPES, 50% Glycerol, 0.87% Sodium chloride, 0.01% BSA
精製度	Immunogen affinity purified
特記事項 (精製)	Prepared from rabbit serum by affinity purification via sequential chromatography on phospho and dephosphopeptide affinity columns.
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

## アプリケーション

**The Abpromise guarantee** Abpromise保証は、次のテスト済みアプリケーションにおけるab30562の使用に適用されます  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		Use at an assay dependent concentration.

## ターゲット情報

機能	Receptor for members of the ephrin-B family. Binds to ephrin-B1, -B2 and -B3. Binding with the guidance cue ephrin-B2 at the optic chiasm midline redirect ventrotemporal (VT) retinal ganglion cells (RGCs) axons ipsilaterally. May be involved in cell-cell interactions in the nervous system.
組織特異性	Preferentially expressed in brain.
配列類似性	Belongs to the protein kinase superfamily. Tyr protein kinase family. Ephrin receptor subfamily. Contains 2 fibronectin type-III domains. Contains 1 protein kinase domain. Contains 1 SAM (sterile alpha motif) domain.
細胞内局在	Membrane.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

## Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors