

Anti-Erk1 (pT202/pY204) + Erk2 (pT185/pY187) antibody [E337] ab32538

リコンビナント **RabMAb**

29 References **画像数 6**

製品の概要

製品名	Anti-Erk1 (pT202/pY204) + Erk2 (pT185/pY187) antibody [E337]
製品の詳細	Rabbit monoclonal [E337] to Erk1 (pT202/pY204) + Erk2 (pT185/pY187)
由来種	Rabbit
特異性	The antibody detects ERK1 phosphorylated on Threonine 202 and Tyrosine 204 and ERK2 phosphorylated on Threonine 185 and Tyrosine 187.
アプリケーション	適用あり: WB, IHC-P, Flow Cyt (Intra), ICC/IF 適用なし: IP
種交差性	交差種: Human
免疫原	Synthetic peptide within Human Erk1 (pT202/pY204) + Erk2 (pT185/pY187). The exact sequence is proprietary. (Peptide available as ab205613)
ポジティブ・コントロール	WB: Serum starved A431 cell lysate treated with EGF. IHC-P: Human thyroid gland cancer tissue. ICC/IF: A431 cells +- EGF.
特記事項	This product is a recombinant monoclonal antibody, which offers several advantages including: <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production For more information see here . Our RabMAb [®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents . Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

バッファー	pH: 7.2 Preservative: 0.01% Sodium azide Constituents: 50% Glycerol (glycerin, glycerine), 0.05% BSA
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	E337
アイソタイプ	IgG


アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab32538の使用に適用されます**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		1/500 - 1/10000. Detects a band of approximately 42, 44 kDa (predicted molecular weight: 42 , 44 kDa).
IHC-P		1/50. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
Flow Cyt (Intra)		1/1000. ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
ICC/IF		1/200 - 1/250.

追加情報 Is unsuitable for IP.

画像




Immunohistochemical analysis of paraffin-embedded human thyroid gland cancer using anti-ERK1(pT202/pY204)/ERK2(pT185/pY187) (ab32538) at dilution of 1:50.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.


Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Erk1 (pT202/pY204) + Erk2 (pT185/pY187) antibody [E337] (ab32538)



Immunocytochemistry/ Immunofluorescence - Anti-Erk1 (pT202/pY204) + Erk2 (pT185/pY187) antibody [E337] (ab32538)

Immunocytochemistry/Immunofluorescence analysis of 4% paraformaldehyde A431+EGF(100ng/ml,5min) labelling Erk1 (pT202/pY204) + Erk2 (pT185/pY187) with ab32538 at dilution of 1/200. The secondary antibody used was Alexa Fluor® 488 Goat-Anti-Rabbit IgG ([ab150077](#)) at dilution of 1/400. The counter stain was done with DAPI (blue).



Western blot - Anti-Erk1 (pT202/pY204) + Erk2 (pT185/pY187) antibody [E337] (ab32538)

Lane 1 : Anti-Erk1 (pT202/pY204) + Erk2 (pT185/pY187) antibody [E337] (ab32538) at 1/500 dilution

Lane 2 : Anti-Erk1 (pT202/pY204) + Erk2 (pT185/pY187) antibody [E337] (ab32538) at 1/2000 dilution

Lane 3 : Anti-Erk1 (pT202/pY204) + Erk2 (pT185/pY187) antibody [E337] (ab32538) at 1/10000 dilution

All lanes : A431 treated with EGF for 10 minutes

Lysates/proteins at 0.1 µg/ml per lane.

Secondary


All lanes : Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 42 , 44 kDa

Observed band size: 42.44 kDa


Exposure time: 3 minutes

First-antibody diluted with 1% BSA.



THP1 cells were incubated at 37°C for 3 minutes with vehicle control (0 μM) and different concentrations of ZK 756326 (**ab120814**). Increased expression of ERK1 (phospho T202 + Y204) + ERK2 (phospho T185 + Y187) (ab32538) in THP1 cells correlates with an increase in ZK 756326 concentration, as described in literature.

Whole cell lysates were prepared with RIPA buffer (containing protease inhibitors and sodium orthovanadate), 10 μg of each were loaded on the gel and the WB was run under reducing conditions. After transfer the membrane was blocked for an hour using 5% BSA before being incubated with ab32538 at 1/500 dilution and **ab17942** at 1 μg/ml overnight at 4°C. Antibody binding was detected using an anti-rabbit antibody conjugated to HRP (**ab97051**) at 1/10000 dilution and visualised using ECL development solution.



Overlay histogram showing HeLa cells stained with ab32538 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab32538, 1/1000 dilution) for 30 min at 22°C. The secondary antibody used was a goat anti-rabbit DyLight® 488 (IgG; H+L) (**ab96899**) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (1 μg/1x10⁶ cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-Erk1 (pT202/pY204) + Erk2 (pT185/pY187)
antibody [E337] (ab32538)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors