


# Anti-FAK antibody [EP695Y] - Low endotoxin, Azide free ab219363

KO 評価済 リコンビナント RabMAb

25 References 画像数 5

### 製品の概要

製品名	Anti-FAK antibody [EP695Y] - Low endotoxin, Azide free
製品の詳細	Rabbit monoclonal [EP695Y] to FAK - Low endotoxin, Azide free
由来種	Rabbit
特異性	<p>ab219363 recognises Focal adhesion kinase (FAK).</p> <p>The mouse and rat recommendation is based on the WB results. We do not guarantee IHC-P for mouse and rat.</p>
アプリケーション	適用あり: WB, IHC-P
種交差性	<p>交差種: Mouse, Rat, Human</p> <p>交差が予測される動物種: Cow </p>
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: NIH/3T3, Rat brain, HeLa, K-562. A431 and HEK-293T lysates IHC-P: human hepatocellular carcinoma
特記事項	<p>ab219363 is the carrier-free version of <a href="#">ab40794</a>.</p> <p>Our <b>carrier-free</b> antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our <b>conjugation kits</b> for antibody conjugates that are ready-to-use in as little as 20 minutes with &lt;1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar<sup>®</sup> Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar<sup>®</sup> is a trademark of Fluidigm Canada Inc.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> </ul>

- Animal-free production

For more information [see here](#).

Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to [RabMAb<sup>®</sup> patents](#).

Our **Low endotoxin, azide-free formats** have low endotoxin level ( $\leq 1$  EU/ml, determined by the LAL assay) and are free from azide, to achieve consistent experimental results in functional assays.

## 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C. Do Not Freeze.
バッファー	pH: 7.20 Constituent: PBS
キャリア・フリー	はい
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EP695Y
アイソタイプ	IgG

## アプリケーション

**The Abpromise guarantee**      **Abpromise保証は、** 次のテスト済みアプリケーションにおけるab219363の使用に適用されます  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。


アプリケーション	Abreviews	特記事項
WB		Use at an assay dependent concentration. Detects a band of approximately 125 kDa (predicted molecular weight: 119 kDa).
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol. The mouse, rat and cow recommendation is based on the WB results. We do not guarantee IHC-P for mouse, rat and cow. See <a href="#">IHC antigen retrieval protocols</a> .

## ターゲット情報

機能	Non-receptor protein-tyrosine kinase implicated in signaling pathways involved in cell motility, proliferation and apoptosis. Activated by tyrosine-phosphorylation in response to either integrin clustering induced by cell adhesion or antibody cross-linking, or via G-protein coupled receptor (GPCR) occupancy by ligands such as bombesin or lysophosphatidic acid, or via LDL receptor occupancy. Microtubule-induced dephosphorylation at Tyr-397 is crucial for the induction of focal adhesion disassembly. Plays a potential role in oncogenic transformations resulting in increased
----	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

	kinase activity.
<b>組織特異性</b>	Expressed in all organs tested, in lymphoid cell lines, but most abundantly in brain.
<b>配列類似性</b>	Belongs to the protein kinase superfamily. Tyr protein kinase family. FAK subfamily. Contains 1 FERM domain. Contains 1 protein kinase domain.
<b>ドメイン</b>	The first Pro-rich domain interacts with the SH3 domain of CRK-associated substrate (BCAR1) and CASL. The carboxy-terminal region is the site of focal adhesion targeting (FAT) sequence which mediates the localization of FAK1 to focal adhesions.
<b>翻訳後修飾</b>	Phosphorylated on 6 tyrosine residues upon activation. Microtubule-induced dephosphorylation at Tyr-397 could be catalyzed by PTPN11 and regulated by ZFYVE21. Dephosphorylated by PTPN11 upon EPHA2 activation by its ligand EFNA1.
<b>細胞内局在</b>	Cell junction > focal adhesion. Cell membrane. Constituent of focal adhesions.

## 画像



**All lanes :** Anti-FAK antibody [EP695Y] ([ab40794](#)) at 1/1000 dilution

**Lane 1 :** HeLa cell lysate

**Lane 2 :** A431 cell lysate

**Lane 3 :** Wild-type HEK-293T cell lysate

**Lane 4 :** PTK2 knockout HEK-293T cell lysate

Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.

**Predicted band size:** 119 kDa


**Observed band size:** 119 kDa

This data was developed using the same antibody clone in a different buffer formulation ([ab40794](#)).

**Lanes 1 - 4:** Merged signal (red and green). Green - [ab40794](#) observed at 119 kDa. Red - loading control, [ab8245](#) observed at 37 kDa.


[ab40794](#) was shown to react with FAK in wild-type HEK-293T cells. Loss of signal was observed when knockout cell line [ab255421](#) (knockout cell lysate [ab263766](#)) was used. Wild-type and FAK knockout samples were subjected to SDS-PAGE. [ab40794](#) and Anti-GAPDH antibody [6C5] - Loading Control

(**ab8245**) were incubated overnight at 4°C at 1 in 1000 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (**ab216776**) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-FAK antibody [EP695Y] - Low endotoxin, Azide free (ab219363)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human hepatocellular carcinoma tissue sections labeling FAK with purified **ab40794** at 1:250 dilution (2.32 µg/ml). Heat mediated antigen retrieval was performed using **ab93684** (Tris/EDTA buffer, pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-FAK antibody [EP695Y] - Low endotoxin, Azide free (ab219363)

Immunohistochemical analysis of paraffin-embedded human hepatocellular carcinoma using **ab40794**. This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab40794**).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-FAK antibody [EP695Y] -

Low endotoxin, Azide free (ab219363)

This image is courtesy of an Abreview submitted by Carl Hobbs.

The image shows FAK antibody (**ab40794**) in human spleen tissue. Clear cytoplasmic positivity in a subset of germinal centre cells. There is intense positivity of the serum in the blood vessels. Endogenous peroxidases were blocked using 2% H<sub>2</sub>O<sub>2</sub>, for 15 minutes.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab40794**).

### Why choose a recombinant antibody?



**Research with confidence**  
Consistent and reproducible results



**Long-term and scalable supply**  
Recombinant technology



**Success from the first experiment**  
Confirmed specificity



**Ethical standards compliant**  
Animal-free production

Anti-FAK antibody [EP695Y] - Low endotoxin, Azide free (ab219363)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet

- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

#### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors