

Anti-FANCC antibody ab5065

★★★★☆ [6 Abreviews](#) [2 References](#) [画像数 2](#)

製品の概要

製品名	Anti-FANCC antibody
製品の詳細	Rabbit polyclonal to FANCC
由来種	Rabbit
特異性	Detects a band at 60kDa in Hela cell lysate corresponding to FANCC.
アプリケーション	適用あり: ICC/IF, WB
種交差性	交差種: Human 非交差種: Mouse
免疫原	Synthetic peptide: CLINKEPQNSGSKLNS , corresponding to amino acids 96 - 112 of Human FANCC. Run BLAST with Run BLAST with

特記事項

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
バッファー	Preservative: 0.01% Sodium azide Constituents: 0.42% Potassium phosphate, 0.878% Sodium chloride
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション


The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab5065の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ICC/IF		Use a concentration of 5 µg/ml.
WB	★★★★★ (6)	1/500 - 1/2000. Detects a band of approximately 60 kDa (predicted molecular weight: 63 kDa).

ターゲット情報

機能	DNA repair protein that may operate in a postreplication repair or a cell cycle checkpoint function. May be implicated in interstrand DNA cross-link repair and in the maintenance of normal chromosome stability. Upon IFNG induction, may facilitate STAT1 activation by recruiting STAT1 to IFNGR1.
組織特異性	Ubiquitous.
関連疾患	Defects in FANCC are the cause of Fanconi anemia complementation group C (FANCC) [MIM:227645]. A disorder affecting all bone marrow elements and resulting in anemia, leukopenia and thrombopenia. It is associated with cardiac, renal and limb malformations, dermal pigmentary changes, and a predisposition to the development of malignancies. At the cellular level it is associated with hypersensitivity to DNA-damaging agents, chromosomal instability (increased chromosome breakage) and defective DNA repair.
発生段階	Expression increases during S phase, is maximal at the G2/M transition, and declines during M phase (at protein level).
細胞内局在	Nucleus. Cytoplasm. The major form is nuclear. The minor form is cytoplasmic.

画像



Anti-FANCC antibody (ab5065) at 1/1500 dilution + HeLa whole cell lysate at 35 µg

Secondary


IRDye 800 conjugated Goat anti-Rabbit IgG [H&L] at 1/10000 dilution

Predicted band size: 63 kDa

Observed band size: 60 kDa

Additional bands at: 33 kDa. We are unsure as to the identity of these extra bands.

Primary antibody incubated with PBS supplemented with 1% normal goat serum and 0.1% milk.



Immunocytochemistry/ Immunofluorescence - Anti-FANCC antibody (ab5065)

ICC/IF image of ab5065 stained MCF7 cells. The cells were 4% formaldehyde fixed (10 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab5065, 5µg/ml) overnight at +4°C. The secondary antibody (green) was **ab96899**, DyLight® 488 goat anti-rabbit IgG (H+L) used at a 1/250 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors