

Anti-FBP1 antibody [EPR4620] ab109732

リコンビナント **RabMAb**

★★★★★ **1 Abreviews** **14 References** 画像数 8

製品の概要

製品名	Anti-FBP1 antibody [EPR4620]
製品の詳細	Rabbit monoclonal [EPR4620] to FBP1
由来種	Rabbit
アプリケーション	適用あり: Flow Cyt (Intra), ICC/IF, WB 適用なし: IHC-P
種交差性	交差種: Mouse, Rat, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: Human stomach, Human fetal liver, Human kidney, Mouse liver, Mouse kidney, Rat liver, and MCF7 lysates and DDDK-tagged FBP1 human full-length recombinant protein. ICC/IF: HeLa cell line Flow Cyt (intra): MCF7 cells.
特記事項	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.
バッファー	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: PBS, 0.05% BSA, 40% Glycerol (glycerin, glycerine)
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EPR4620

アイソタイプIgG

アプリケーション

The Abpromise guarantee Abpromise保証は、 次のテスト済みアプリケーションにおけるab109732の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt (Intra)		1/400.
ICC/IF		Use a concentration of 10 µg/ml.
WB	★★★★★ (1)	1/1000 - 1/10000. Detects a band of approximately 36 kDa (predicted molecular weight: 37 kDa).

追加情報Is unsuitable for IHC-P.


ターゲット情報

パスウェイCarbohydrate biosynthesis; gluconeogenesis.

関連疾患Defects in FBP1 are the cause of fructose-1,6-bisphosphatase deficiency (FBPD) [MIM:229700]. FBPD is inherited as an autosomal recessive disorder mainly in the liver and causes life-threatening episodes of hypoglycemia and metabolic acidosis (lactacidemia) in newborn infants or young children.

配列類似性Belongs to the FBPase class 1 family.

画像



All lanes : Anti-FBP1 antibody [EPR4620] (ab109732) at 1/1000 dilution

Lane 1 : DDDK-tagged FBP1 human full-length recombinant protein, 15 ng

Lane 2 : DDDK-tagged FBP12 human full-length recombinant protein, 15 ng

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution


Predicted band size: 37 kDa

Observed band size: 37 kDa

Exposure time: 5 seconds


Blocking buffer and concentration: 5% NFDM/TBST.

Diluting buffer and concentration: 5% NFDM/TBST.



Immunocytochemistry/ Immunofluorescence - Anti-FBP1 antibody [EPR4620] (ab109732)

Immunocytochemistry/ Immunofluorescence analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling FBP1 with purified ab109732 at 1:50 dilution (7.9 µg/ml). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Cells were counterstained with Ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1:200 (2.5 µg/ml). Goat anti rabbit IgG (Alexa Fluor® 488, [ab150077](#)) was used as the secondary antibody at 1:1000 (2 µg/ml) dilution. DAPI nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



Western blot - Anti-FBP1 antibody [EPR4620] (ab109732)

All lanes : Anti-FBP1 antibody [EPR4620] (ab109732) at 1/1000 dilution (Purified)

Lane 1 : Human kidney lysates

Lane 2 : Mouse liver lysates

Lane 3 : Mouse kidney lysates

Lane 4 : Rat liver lysates


Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

Predicted band size: 37 kDa

Observed band size: 36 kDa



Western blot - Anti-FBP1 antibody [EPR4620]
(ab109732)


Anti-FBP1 antibody [EPR4620] (ab109732) at 1/50000 dilution +
MCF7 (Human breast adenocarcinoma epithelial cell) whole cell
lysates at 15 µg

Secondary

Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution


Predicted band size: 37 kDa

Observed band size: 36 kDa




Flow Cytometry (Intracellular) - Anti-FBP1 antibody
[EPR4620] (ab109732)

Intracellular Flow Cytometry analysis of MCF7 (Human breast
adenocarcinoma epithelial cell) cells labeling FBP1 with purified
ab109732 at 1/400 dilution (1 µg/ml) (red). Cells were fixed with 4%
Paraformaldehyde. A Goat anti rabbit IgG (Alexa Fluor® 488,
[ab150077](#)) secondary antibody was used at 1/2000. Isotype control
- Rabbit monoclonal IgG (Black). Unlabeled control - Cell without
incubation with primary antibody and secondary antibody (Blue).



Immunocytochemistry/ Immunofluorescence - Anti-
FBP1 antibody [EPR4620] (ab109732)

ICC/IF image of ab109732 stained HeLa cells. The cells were 4% formaldehyde fixed (10 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab109732, 10µg/ml) overnight at +4°C. The secondary antibody (green) was **ab96899**, DyLight® 488 goat anti-rabbit IgG (H+L) used at a 1/250 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.



Western blot - Anti-FBP1 antibody [EPR4620]
(ab109732)

All lanes : Anti-FBP1 antibody [EPR4620] (ab109732) at 1/1000 dilution

Lane 1 : Human stomach lysate

Lane 2 : Human fetal liver lysate

Lane 3 : MCF7 lysate

Lysates/proteins at 10 µg per lane.

Predicted band size: 37 kDa

Why choose a
recombinant antibody?



**Research with
confidence**
Consistent and
reproducible results



**Long-term and
scalable supply**
Recombinant
technology



**Success from the
first experiment**
Confirmed
specificity



**Ethical standards
compliant**
Animal-free
production

Anti-FBP1 antibody [EPR4620] (ab109732)

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors