

Anti-FGF17 antibody ab9990

画像数 1

製品の概要

製品名	Anti-FGF17 antibody
製品の詳細	Rabbit polyclonal to FGF17
由来種	Rabbit
アプリケーション	適用あり: Sandwich ELISA
免疫原	Recombinant full length protein corresponding to Human FGF17 aa 1-250. Database link: O60258
ポジティブ・コントロール	 Run BLAST with  Run BLAST with Recombinant human FGF17 protein (ab9991) can be used as a positive control in WB.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Lyophilized:Reconstitute with 200µl of sterile water. Please note that if you receive this product in liquid form it has already been reconstituted as described and no further reconstitution is necessary.
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term.
バッファー	No preservative, sterile filtered
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	unknown
軽鎖の種類	unknown

アプリケーション

The Abpromise guarantee

Abpromise保証は、次のテスト済みアプリケーションにおけるab9990の使用に適用されます


アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Sandwich ELISA		Use at an assay dependent concentration. Can be paired for Sandwich ELISA with Mouse monoclonal [MM0284-5H27] to FGF17 (ab89270) . For sandwich ELISA, use this antibody as Detection at 0.5µg/ml with ab89270 as Capture.

ターゲット情報

機能	Plays an important role in the regulation of embryonic development and as signaling molecule in the induction and patterning of the embryonic brain. Required for normal brain development.
組織特異性	Preferentially expressed in the embryonic brain.
配列類似性	Belongs to the heparin-binding growth factors family.
発生段階	Detected in embryos at E14.5, but not at E10.5 and E19.5. Preferentially expressed in the neuroepithelia of the isthmus and septum of the embryonic brain at E14.5.
細胞内局在	Secreted.

画像



Standard Curve for FGF17 (Analyte: **ab9991**) dilution range 1pg/ml to 1ug/ml using Capture Antibody Mouse monoclonal [MM0284-5H27] to FGF17 (**ab89270**) at 1ug/ml and Detector Antibody Rabbit polyclonal to FGF17 (ab9990) at 0.5ug/ml

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors