

FITC Anti-CD84 antibody [CD84.1.21] ab33217

製品の概要

製品名	FITC Anti-CD84 antibody [CD84.1.21]
製品の詳細	FITC Mouse monoclonal [CD84.1.21] to CD84
由来種	Mouse
標識	FITC. Ex: 493nm, Em: 528nm
特異性	Recognizes CD84 cell surface antigen, a highly glycosylated cell surface antigen of 64-82kD.
アプリケーション	適用あり: Flow Cyt
種交差性	交差種: Human
免疫原	Tissue, cells or virus corresponding to CD84. Mouse pre-B cell line 300.19 transfected with Human CD84 cDNA.
ポジティブ・コントロール	Whole blood
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C.
バッファー	Preservative: 0.09% Sodium azide Constituents: PBS, 1% BSA
精製度	Protein G purified
特記事項 (精製)	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
ポリ/モノ	モノクローナル
クローン名	CD84.1.21
ミエローマ	NS1
アイソタイプ	IgG1

アプリケーション

The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab33217の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt		Use 10µl for 10 ⁶ cells. ab91356 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody. Use 10µl for 10 ⁶ cells or 100µl whole blood.

ターゲット情報

機能	Plays a role as adhesion receptor functioning by homophilic interactions and by clustering. Recruits SH2 domain-containing proteins SH2D1A/SAP. Increases proliferative responses of activated T-cells and SH2D1A/SAP does not seem to be required for this process. Homophilic interactions enhance interferon gamma/IFNG secretion in lymphocytes and induce platelet stimulation via a SH2D1A/SAP-dependent pathway. May serve as a marker for hematopoietic progenitor cells.
組織特異性	Predominantly expressed in hematopoietic tissues, such as lymph node, spleen and peripheral leukocytes. Expressed in macrophages, B-cells, monocytes, platelets, thymocytes, T-cells and dendritic cells. Highly expressed in memory T-cells.
配列類似性	Contains 1 Ig-like C2-type (immunoglobulin-like) domain.
発生段階	Expression is slightly increased in naive B-cells after the first division. By contrast, expression on memory B-cells decreased with each successive division.
ドメイン	ITSM (immunoreceptor tyrosine-based switch motif) motif is a cytoplasmic motif which may bind SH2D1A.
翻訳後修飾	Phosphorylated by tyrosine-protein kinase LCK on tyrosine residues following ligation induced by agonist monoclonal antibody. The association with SH2D1A/SAP is dependent of tyrosines phosphorylation of its cytoplasmic domain Phosphorylated on Tyr-296 and Tyr-316 following platelet aggregation. N-glycosylated.
細胞内局在	Cell membrane.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish

- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors