

FITC Anti-HIV1 p24 antibody ab20569

1 References

製品の概要

製品名	FITC Anti-HIV1 p24 antibody
製品の詳細	FITC Goat polyclonal to HIV1 p24
由来種	Goat
標識	FITC. Ex: 493nm, Em: 528nm
アプリケーション	適用あり: ELISA, WB
種交差性	交差種: Human immunodeficiency virus 非交差種: Cow, Human
免疫原	Full length native protein (purified) for strain III _B .
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C.
バッファー	Preservative: 0.1% Sodium azide Constituents: 0.0268% PBS, 1% BSA
精製度	IgG fraction
特記事項 (精製)	Purified IgG fraction covalently coupled with high purity Isomer I of fluorescein isothiocyanate. Care is taken to ensure complete removal of any free fluorescein from the final product.
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab20569の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ELISA		
WB		

追加情報 ELISA: Use at an assay dependent dilution. WB: Use at an assay dependent dilution. Dilution optimised using Chromogenic detection. Direct FA staining of target antigens in a permissive tissue culture system. Working dilution must be determined by the user but a starting range of 1:10 - 1:50 is suggested. Acetone fixation of the antigen source is recommended prior to staining. Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user.

ターゲット情報

関連性 HIV1 performs highly complex orchestrated tasks during the assembly, budding, maturation and infection stages of the viral replication cycle. During viral assembly, the proteins form membrane associations and self-associations that ultimately result in budding of an immature virion from the infected cell. Gag precursors also function during viral assembly to selectively bind and package two plus strands of genomic RNA. Capsid protein p24 probably forms the conical core of the virus that encapsulates the genomic RNA-nucleocapsid complex.

細胞内局在 Membrane

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors