abcam

Product datasheet

FITC Anti-HLA A2 antibody [BB7.2] ab27728

8 References 画像数 1

製品の概要

製品名 FITC Anti-HLA A2 antibody [BB7.2]

製品の詳細 FITC Mouse monoclonal [BB7.2] to HLA A2

由来種 Mouse

標識 FITC. Ex: 493nm, Em: 528nm

特異性 Ab27728 recognises the human HLA A2 histocompatability antigen.

アプリケーション 適用あり: Flow Cyt

種交差性 交差種: Human

免疫原 Full length protein corresponding to Human HLA A2. Papain solubilized HLA-A2

エピトープ The epitope recognised by this antibody has been studied extensively and would appear to

include the carboxy terminus of the alpha 2 helix and a turn on one of the underlying beta strands.

特記事項

The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C.

パッファー Preservative: 0.09% Sodium azide

Constituent: 1% BSA

精製度 Protein G purified

特記事項(精製) Ab27728 was purified from tissue culture supernatant, and prepared by affinity chromatography.

ポリ/モノ モノクローナル

クローン名 BB7.2 ミ**エローマ** NS1

1

アプリケーション

The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab27728の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt		1/10. Use 10 μl of the suggested working dilution to label 106 cells or 100μl whole blood.

ターゲット情報


関連性

Involved in the presentation of foreign antigens to the immune system. Dimer of alpha chain and a beta chain (beta-2-microglobulin). Interacts with human herpesvirus 8 MIR1 protein.

細胞内局在

Membrane; Single-pass type I membrane protein.

画像



Ab27728, undiluted or at a dilution of 1/10, staining HLA A2 in human peripheral blood monocytes by FACS.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery

- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors