


Anti-FOXP1 antibody ab16645

★★★★★ 22 Abreviews 97 References 画像数 3

製品の概要

製品名	Anti-FOXP1 antibody
製品の詳細	Rabbit polyclonal to FOXP1
由来種	Rabbit
特異性	Replenishment batches of our polyclonal antibody, ab16645 are tested in IHC-P. Previous batches were additionally validated in IHC-FoFr and WB. These applications are still expected to work and are covered by our Abpromise guarantee. You may also be interested in our alternative recombinant antibody, ab134055 .
アプリケーション	適用あり: IHC-FoFr, IHC-P, WB
種交差性	交差種: Mouse 交差が予測される動物種: Rat, Chicken, Dog, Human, Xenopus laevis 
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	IHC-p: Mouse brain
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	pH: 7.40 Preservative: 0.02% Sodium azide Constituent: PBS
	Batches of this product that have a concentration < 1mg/ml may have BSA added as a stabilising agent. If you would like information about the formulation of a specific lot, please contact our

	scientific support team who will be happy to help.
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション


The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab16645の使用に適用されます**
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-FoFr		Use at an assay dependent concentration.
IHC-P	★★★★★ (4)	Use a concentration of 0.1 µg/ml. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
WB	★★★★★ (7)	1/5000. Detects a band of approximately 75 kDa (predicted molecular weight: 75 kDa). ab16645 recognises the different isoforms of FOXP1 in Jurkat cell lysate (see review for further details).

ターゲット情報


機能	Transcriptional repressor. It plays an important role in the specification and differentiation of lung epithelium. Can act with CTBP1 to synergistically repress transcription but CTPBP1 is not essential. Essential transcriptional regulator of B cell development.
関連疾患	Note=A chromosomal aberration involving FOXP1 is found in acute lymphoblastic leukemia. Translocation t(9;3)(p13;p14.1) with PAX5. Defects in FOXP1 are the cause of mental retardation with language impairment and autistic features (MRLIAF) [MIM:613670]. It is a developmental disorder characterized by mild to moderate mental retardation, language impairment, and autistic features. Patients show global delay, delayed walking, severely delayed speech development, and behavioral abnormalities, including irritability, hyperactivity, aggression, and stereotypical rigid behaviors.
配列類似性	Contains 1 C2H2-type zinc finger. Contains 1 fork-head DNA-binding domain.
ドメイン	The leucine-zipper is required for dimerization and transcriptional repression.
細胞内局在	Nucleus.

画像



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-FOXP1 antibody (ab16645)

Paraffin-embedded mouse brain tissue stained for FOXP1 using ab16645 in immunohistochemical analysis.



Western blot - Anti-FOXP1 antibody (ab16645)

All lanes : Anti-FOXP1 antibody (ab16645) at 1/5000 dilution

Lane 1 : Lysate from e12.5 whole mouse embryo FoxP1 +/-

Lane 2 : Lysate from e12.5 whole mouse embryo FoxP1 -/-


Lysates/proteins at 30 µg per lane.

Developed using the ECL technique.

Predicted band size: 75 kDa

Observed band size: 65,82 kDa

Additional bands at: 45 kDa (possible cross reactivity), 50 kDa (possible cross reactivity), 85 kDa (possible cross reactivity)



Immunohistochemistry (PFA perfusion fixed frozen sections) - Anti-FXP1 antibody (ab16645)

This image is courtesy of Shunyou Wang, Ph.D.

ab16645 was tested in IHC of formalin fixed sections from mouse (over two weeks of age) brain. Antigen retrieval was in citrate buffer and blocking was in 1% BSA. The primary antibody was incubated at a dilution of 1/200 overnight at 4 degrees celsius. The secondary antibody was a biotin conjugated anti-rabbit IgG.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors