abcam

Product datasheet

Anti-FSTL1/FRP antibody ab71548

★★★★★ 1 Abreviews 7 References

製品の概要

特記事項

製品名 Anti-FSTL1/FRP antibody

製品の詳細 Rabbit polyclonal to FSTL1/FRP

由来種 Rabbit

アプリケーション 適用あり: IHC-P, WB

種交差性 交差種: Human

交差が予測される動物種: Mouse, Rat 🔷

免疫原 Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

We are constantly working hard to ensure we provide our customers with best in class antibodies. As a result, we are pleased to offer this antibody in a purified format as of 19th February 2018. The following lots is/are still unpurified and still in stock as of 19th February 2018 - GR200980-7, GR200980-9, GR200980-10. Lot numbers other than GR200980-7, GR200980-9, GR200980-10 will be purified. Please note that the dilutions may need to be adjusted accordingly. Purified antibodies have the advantage of being enriched for the fraction of immunoglobulin that specifically reacts with the target antigen and for having a reduction of serum proteins.

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.

バッファー Constituents: 0.92% Glycerol, 1.21% Tris, 2% Sucrose

精製度 Protein A purified

ポリ/モノ ポリクローナル

アイソタイプ lgG

1

The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab71548の使用に適用されますアプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-P		Use at an assay dependent concentration.
WB	★★★ ☆☆ <u>(1)</u>	Use at an assay dependent concentration.

ターゲット情報

機能 May modulate the action of some growth factors on cell proliferation and differentiation. Binds

heparin.

組織特異性 Overexpressed in synovial tissues from rheumatoid arthritis.

配列類似性Contains 2 EF-hand domains.
Contains 1 follistatin-like domain.

Contains 1 Kazal-like domain.

Contains 1 WFC domain.

細胞内局在 Secreted.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors