


Anti-FYTTD1 antibody ab125082

★★★★★ [1 Abreviews](#) [画像数 2](#)

製品の概要

| | |
|--------------|---|
| 製品名 | Anti-FYTTD1 antibody |
| 製品の詳細 | Rabbit polyclonal to FYTTD1 |
| 由来種 | Rabbit |
| アプリケーション | 適用あり: WB, IP |
| 種交差性 | 交差種: Mouse, Human 交差が予測される動物種: Rat, Horse, Chimpanzee, Rhesus monkey, Gorilla, Orangutan, Bat, Elephant  |
| 免疫原 | Synthetic peptide, corresponding to a region within amino acids 1-50 of Human FYTTD1 (NP_115664.2). |
| ポジティブ・コントロール | 293T, Jurkat, HeLa and NIH3T3 whole cell lysates. |
| 特記事項 | <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p> |

製品の特性

| | |
|-----------|--|
| 製品の状態 | Liquid |
| 保存方法 | Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles. |
| バッファー | <p>pH: 7</p> <p>Preservative: 0.09% Sodium azide</p> <p>Constituent: 99% Tris citrate/phosphate</p> <p>pH 7 to 8</p> |
| 精製度 | Immunogen affinity purified |
| 特記事項 (精製) | ab125082 is affinity purified using an epitope specific to FYTTD1 immobilized on solid support. |
| ポリ/モノ | ポリクローナル |

アプリケーション


The Abpromise guarantee **Abpromise保証は、** 次のテスト済みアプリケーションにおけるab125082の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

| アプリケーション | Abreviews | 特記事項 |
|----------|-----------|--|
| WB | ★★★★★ (1) | 1/1000 - 1/5000. Predicted molecular weight: 36 kDa. |
| IP | | Use at 2-10 µg/mg of lysate. |

ターゲット情報

| | |
|-------|---|
| 機能 | Required for mRNA export from the nucleus to the cytoplasm. Acts as an adapter that uses the DDX39B/UAP56-NFX1 pathway to ensure efficient mRNA export and delivering to the nuclear pore. Associates with spliced and unspliced mRNAs simultaneously with THOC4. |
| 配列類似性 | Belongs to the UIF family. |
| 細胞内局在 | Nucleus > nucleoplasm. Nucleus speckle. |

画像




All lanes : Anti-FYTTD1 antibody (ab125082) at 0.4 µg/ml

- Lane 1 :** 293T whole cell lysate at 50 µg
- Lane 2 :** 293T whole cell lysate at 15 µg
- Lane 3 :** Jurkat whole cell lysate at 50 µg
- Lane 4 :** HeLa cell lysate at 50 µg
- Lane 5 :** NIH3T3 whole cell lysate at 50 µg

Developed using the ECL technique.

Predicted band size: 36 kDa

Exposure time: 30 seconds



Detection of FYTDD1 by Western Blot of Immunoprecipitate.
ab125082 at 1 µg/ml staining FYTDD1 in 293T whole cell lysate
immunoprecipitated using ab125082 at 6 µg/mg lysate (1 mg/IP;
20% of IP loaded/lane).
Detection: Chemiluminescence with exposure time of 30 seconds.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors