





Anti-GARNL3 antibody ab121712

画像数 2

製品の概要

製品名	Anti-GARNL3 antibody
製品の詳細	Rabbit polyclonal to GARNL3
由来種	Rabbit
アプリケーション	適用あり: WB, IHC-P
種交差性	交差種: Human
免疫原	antigen sequence, corresponding to amino acids 503-598 (LVSTDAGVLL VDDDLPSVPV FDRTLVPVKQM HVLETLDLLV LRADKGGDAR LFFVRLSALQ KGLEGKQAGK SRSDCRENKL EKTGCHLYA INTHHS) of Human GARNL3.  Run BLAST with   Run BLAST with 
ポジティブ・コントロール	Human smooth muscle tissue.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
バッファー	pH: 7.20 Preservative: 0.02% Sodium azide Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine)
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee **Abpromise保証は、**次のテスト済みアプリケーションにおけるab121712の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。


アプリケーション	Abreviews	特記事項
WB		Use a concentration of 0.04 - 0.4 µg/ml.
IHC-P		1/50 - 1/200. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

ターゲット情報

配列類似性


Belongs to the GARNL3 family.
Contains 1 CNH domain.
Contains 1 Rap-GAP domain.

画像



ab121712, at 1/50 dilution, staining GARNL3 in paraffin-embedded Human smooth muscle tissue, by Immunohistochemistry.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-GARNL3 antibody (ab121712)



Western blot - Anti-GARNL3 antibody (ab121712)

Lane 1: Marker [kDa] 250; 130; 95; 72; 55; 36; 28; 17; 10
 Lane 2: Negative control (vector only transfected HEK293T lysate)
 Lane 3: Over-expression Lysate (Co-expressed with a C-terminal myc-DDK tag (~3.1 kDa) in mammalian HEK293T cells. Lane 1: Marker [kDa] 250; 130; 95; 72; 55; 36; 28; 17; 10
 Lane 2: Negative control (vector only transfected HEK293T lysate)
 Lane 3: Over-expression Lysate (Co-expressed with a C-terminal myc-DDK tag (~3.1 kDa) in mammalian HEK293T cells.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors