

Anti-GFAP antibody ab134436

★★★★★ [3 Abreviews](#) [17 References](#) [画像数 2](#)

製品の概要

製品名	Anti-GFAP antibody
製品の詳細	Chicken polyclonal to GFAP
由来種	Chicken
アプリケーション	適用あり: ICC/IF
種交差性	交差種: Rat
免疫原	<p>This product was produced with the following immunogens:</p> <p>Recombinant full length protein corresponding to Cow GFAP. Recombinant full length protein followed by boosts of native GFAP protein purified from bovine spinal cords</p> <p>Full length native protein (purified) corresponding to Cow GFAP. Recombinant full length protein corresponding to GFAP. Recombinant full length protein followed by boosts of native GFAP protein purified from bovine spinal cords</p>
特記事項	<p>Do not freeze this antibody unless you want to store them for longer periods of time. Note, however, that each time an antibody preparation is frozen, about half its binding activity is lost.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	<p>pH: 7</p> <p>Preservative: 0.02% Sodium azide</p> <p>Constituents: 99% PBS, 0.5% BSA</p>
精製度	IgY fraction
特記事項 (精製)	After repeated injections, immune eggs were collected, the IgY fractions were purified from the

yolks, and the IgY concentration adjusted to 20 mg/ml. This preparation was then diluted 1:10 with PBS containing bovine serum albumin as a carrier. Finally, the antibody preparation was filter-sterilized.

ポリ/モノ
アイソタイプ

ポリクローナル
IgY

アプリケーション


The Abpromise guarantee Abpromise保証は、 次のテスト済みアプリケーションにおけるab134436の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
ICC/IF	★★★★★ (1)	1/2000 - 1/5000. Use 2% paraformaldehyde-fixed cells.

ターゲット情報

機能	GFAP, a class-III intermediate filament, is a cell-specific marker that, during the development of the central nervous system, distinguishes astrocytes from other glial cells.
組織特異性	Expressed in cells lacking fibronectin.
関連疾患	Defects in GFAP are a cause of Alexander disease (ALEXD) [MIM:203450]. Alexander disease is a rare disorder of the central nervous system. It is a progressive leukoencephalopathy whose hallmark is the widespread accumulation of Rosenthal fibers which are cytoplasmic inclusions in astrocytes. The most common form affects infants and young children, and is characterized by progressive failure of central myelination, usually leading to death usually within the first decade. Infants with Alexander disease develop a leukoencephalopathy with macrocephaly, seizures, and psychomotor retardation. Patients with juvenile or adult forms typically experience ataxia, bulbar signs and spasticity, and a more slowly progressive course.
配列類似性	Belongs to the intermediate filament family.
翻訳後修飾	Phosphorylated by PKN1.
細胞内局在	Cytoplasm. Associated with intermediate filaments.


画像



Immunofluorescent analysis of dissociated culture of rat cerebral cortical neurons and astrocytes labelling GFAP with ab134436 at 1/2000 dilution (red staining). The nuclei of some neuronal cells (blue staining) are visible.

Immunocytochemistry/ Immunofluorescence - Anti-GFAP antibody (ab134436)

Picture courtesy of Dr. Gerry Shaw, University of Florida



ICC image of ab134436 stained primary neuron-glia rat cells. The cells were fixed with 3.7% formaldehyde in PBS (5 min), permeabilized with cold methanol and then blocked with 2% goat serum for 1h. The cells were then incubated with ab134436 (1/5000) overnight at +4°C followed by a further incubation at room temperature with Goat Anti-chicken IgG H&L (AlexaFluor® 488) at 1/2000 dilution (shown in red) for 1hr. Nuclear DNA was labelled in blue with DAPI.

Immunocytochemistry/ Immunofluorescence - Anti-GFAP antibody (ab134436)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors