

Anti-GFPT1 antibody [EPR4854] ab125069

リコンビナント **RabMAb**

★★★★★ **2 Abreviews** **22 References** 画像数 7

製品の概要

製品名	Anti-GFPT1 antibody [EPR4854]
製品の詳細	Rabbit monoclonal [EPR4854] to GFPT1
由来種	Rabbit
特異性	The mouse and rat recommendation is based on the WB results. We do not guarantee IHC-P for mouse and rat.
アプリケーション	適用あり: WB, IP, IHC-P 適用なし: Flow Cyt or ICC/IF
種交差性	交差種: Mouse, Rat, Human
免疫原	Synthetic peptide within Human GFPT1 aa 600-700. The exact sequence is proprietary.
ポジティブ・コントロール	IHC-P: Human testis, and Human breast carcinoma tissue; WB: Human placenta lysate, MCF7, C6, MEF, 293T, JAR and HeLa cell line lysates, Mouse heart and Mouse cerebral cortex, Rat heart tissue lysate. IP: Jurkat cells.
特記事項	This product is a recombinant monoclonal antibody, which offers several advantages including: <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production For more information see here . Our RabMAb [®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents .

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.
バッファー	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol, 0.05% BSA
精製度	Protein A purified
ポリ/モノ	モノクローナル

クローン名	EPR4854
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee **Abpromise保証は、**次のテスト済みアプリケーションにおけるab125069の使用に適用されます
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。


アプリケーション	Abreviews	特記事項
WB	★★★★★ (2)	1/200. Detects a band of approximately 79 kDa (predicted molecular weight: 79 kDa). For unpurified use at 1/1000 - 1/1000.
IP		1/10 - 1/100.
IHC-P		1/1000. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol. For unpurified use at 1/100 - 1/250. See IHC antigen retrieval protocols . The mouse and rat recommendation is based on the WB results. We do not guarantee IHC-P for mouse and rat.

追加情報 Is unsuitable for Flow Cyt or ICC/IF.

ターゲット情報


機能	Controls the flux of glucose into the hexosamine pathway. Most likely involved in regulating the availability of precursors for N- and O-linked glycosylation of proteins.
組織特異性	Isoform 1 is predominantly expressed in skeletal muscle. Not expressed in brain. Seems to be selectively expressed in striated muscle.
パスウェイ	Nucleotide-sugar biosynthesis; UDP-N-acetyl-alpha-D-glucosamine biosynthesis; alpha-D-glucosamine 6-phosphate from D-fructose 6-phosphate: step 1/1.
関連疾患	Defects in GFPT1 are the cause of limb-girdle myasthenia with tubular aggregates (LGMTA) [MIM:610542]. A congenital myasthenic syndrome characterized by onset of proximal muscle weakness in the first decade. Individuals with this condition have a recognizable pattern of weakness of shoulder and pelvic girdle muscles, and sparing of ocular or facial muscles. EMG classically shows a decremental response to repeated nerve stimulation, a sign of neuromuscular junction dysfunction. Affected individuals show a favorable response to acetylcholinesterase (AChE) inhibitors.
配列類似性	Contains 1 glutamine amidotransferase type-2 domain. Contains 2 SIS domains.

画像




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-GFPT1 antibody
[EPR4854] (ab125069)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human breast carcinoma tissue sections labeling GFPT1 with Purified ab125069 at 1:1000 dilution (2.3 µg/ml). Heat mediated antigen retrieval was performed using **ab93684** (Tris/EDTA buffer, pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.



Immunoprecipitation - Anti-GFPT1 antibody
[EPR4854] (ab125069)

ab125069 (purified) at 1:100 dilution (2µg) immunoprecipitating GFPT1 in Jurkat whole cell lysate.
Lane 1 (input): Jurkat (Human T cell leukemia T lymphocyte) whole cell lysate 10µg
Lane 2 (+): ab125069 & Jurkat whole cell lysate
Lane 3 (-): Rabbit monoclonal IgG (**ab172730**) instead of ab125069 in Jurkat whole cell lysate
For western blotting, VeriBlot for IP Detection Reagent (HRP) (**ab131366**) was used for detection at 1:1000 dilution.
Blocking and diluting buffer: 5% NFDM/TBST.



Western blot - Anti-GFPT1 antibody [EPR4854]
(ab125069)


Anti-GFPT1 antibody [EPR4854] (ab125069) at 1/200 dilution
(Purified) + C6 (Rat glial tumor glial cell) whole cell lysates at 15 µg

Secondary

Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

Predicted band size: 79 kDa

Observed band size: 79 kDa



Western blot - Anti-GFPT1 antibody [EPR4854]
(ab125069)

All lanes : Anti-GFPT1 antibody [EPR4854] (ab125069) at 1/1000 dilution (Purified)

Lane 1 : MCF7 (Human breast adenocarcinoma epithelial cell) whole cell lysates

Lane 2 : Mouse heart lysates

Lane 3 : Mouse cerebral cortex lysates

Lane 4 : MEF (Mouse embryonic fibroblast (immortalized)) whole cell lysates

Lane 5 : Rat heart lysates

Lysates/proteins at 20 µg per lane.


Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

Predicted band size: 79 kDa

Observed band size: 79 kDa


This antibody is not suitable to detect tissue lysate samples.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-GFPT1 antibody [EPR4854] (ab125069)

ab125069, at a 1/100 dilution, staining GFPT1 in paraffin-embedded Human testis tissue by immunohistochemistry.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Western blot - Anti-GFPT1 antibody [EPR4854] (ab125069)

All lanes : Anti-GFPT1 antibody [EPR4854] (ab125069) at 1/1000 dilution

Lane 1 : 293T cell lysate

Lane 2 : JAR cell lysate

Lane 3 : HeLa cell lysate

Lane 4 : Human placenta lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat anti-Rabbit HRP at 1/2000 dilution

Developed using the ECL technique.

Predicted band size: 79 kDa

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-GFPT1 antibody [EPR4854] (ab125069)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors