abcam

Product datasheet

Anti-GST3 / GST pi antibody [EPR8263] ab138491



יוכציטו RabMAb

★★★★★ 1 Abreviews 11 References 画像数9

製品の概要

製品名 Anti-GST3 / GST pi antibody [EPR8263]

製品の詳細 Rabbit monoclonal [EPR8263] to GST3 / GST pi

由来種 Rabbit

アプリケーション 適用あり: Flow Cyt (Intra), WB, IHC-P, ICC/IF

適用なし: IP

種交差性 交差種: Mouse, Rat, Human

免疫原 Synthetic peptide within Human GST3/ GST pi aa 100-200. The exact sequence is proprietary.

ポジティブ・コントロール WB: HAP1, Jurkat, and HEK293 cell lysates; ICC/IF: K562 cells; Flow Cyt (intra): Jurkat; IHC-P:

Rat and mouse stomach, and human tonsil tissue sections; WB: K-562 whole cell lysate, mouse

and rat brain lysates.

特記事項 This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity

- Long-term security of supply

- Animal-free production

For more information see here.


Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at -20°C.

 $K_D = 1.00 \times 10^{-10} M$ 解離定数(KD値)



Learn more about K_D

バッファー pH: 7.20 Preservative: 0.01% Sodium azide

Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA

精製度 Protein A purified

ポリ(モノ モノクローナル **クローン名** EPR8263

アイソタイプ lgG

アプリケーション

The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab138491の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt (Intra)		1/20.
WB		1/1000 - 1/10000. Predicted molecular weight: 23 kDa.
IHC-P	★★★★ (1)	1/100. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol. For unpurified please use at 1/250 - 1/500 dilution
ICC/IF		1/50. For unpurified use at 1/100 - 1/250

追加情報 Is unsuitable for IP.

ターゲット情報

機能 Conjugation of reduced glutathione to a wide number of exogenous and endogenous hydrophobic

electrophiles.

配列類似性 Belongs to the GST superfamily. Pi family.

Contains 1 GST C-terminal domain.
Contains 1 GST N-terminal domain.

画像



Western blot - Anti-GST3 / GST pi antibody [EPR8263] (ab138491)

All lanes: Anti-GST3 / GST pi antibody [EPR8263] (ab138491) at 1/1000 dilution (Purified)

Lane 1: K-562 (Human chronic myelogenous leukemia lymphoblast) whole cell lysate

Lane 2 : Mouse brain lysate


Lane 3: Rat brain lysate

Lysates/proteins at 15 µg per lane.

Secondary


All lanes : Goat Anti-Rabbit lgG H&L (HRP) (<u>ab97051</u>) at 1/20000 dilution

Predicted band size: 23 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-GST3 / GST pi antibody [EPR8263] (ab138491)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of rat stomach tissue sections labeling GST3 / GST pi with Purified ab138491 at 1/100 dilution (0.99 µg/mL). Heat mediated antigen retrieval using Bond™ Epitope Retrieval Solution 1 (pH 6.0). Rabbit specific IHC polymer detection kit HRP/DAB (ab209101) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.



Western blot - Anti-GST3 / GST pi antibody [EPR8263] (ab138491)

Lane 1: Wild type HAP1 whole cell lysate (20 µg)


Lane 2: GST3 knockout HAP1 whole cell lysate (20 µg)

Lane 3: Jurkat whole cell lysate (20 µg)

Lane 4: Hek293 whole cell lysate (20 µg)


Lanes 1 - 4: Merged signal (red and green). Green - ab138491 observed at 23 kDa. Red - loading control, **ab9484**, observed at 37 kDa.

ab138491 was shown to specifically react with GST3 when GST3 knockout samples were used. Wild-type and GST3 knockout samples were subjected to SDS-PAGE. Ab138491 and **ab9484** (Mouse anti GAPDH loading control) were incubated overnight at 4°C at 1000 dilution and 1/10001 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed **ab216773** and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed **ab216776** secondary antibodies at 1/10000 dilution for 1 hour at room temperature before imaging.




Immunocytochemistry/ Immunofluorescence - Anti-GST3 / GST pi antibody [EPR8263] (ab138491)

Immunocytochemistry analysis of K-562 (Human chronic myelogenous leukemia lymphoblast) cells labeling GST3 / GST pi with Purified ab138491 at 1/50 dilution (1.98 μ g/mL). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Cells were counterstained with Ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1/200 dilution (2.5 μ g/mL). Goat anti rabbit lgG (Alexa Fluor® 488, **ab150077**) was used as the secondary antibody at 1/1000 dilution (2 μ g/mL). DAPI (blue) was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.




Flow Cytometry (Intracellular) - Anti-GST3 / GST pi antibody [EPR8263] (ab138491) Intracellular Flow Cytometry analysis of Jurkat (Human T cell leukemia T lymphocyte) cells labeling GST3 / GST pi with Purified ab138491 at 1/20 dilution (10 $\mu g/mL$) (Red). Cells were fixed with 4% Paraformaldehyde and permeabilised with 90% Methanol. A Goat anti rabbit lgG (Alexa Fluor $^{\mbox{\scriptsize (B)}}$ 488, $\mbox{\it ab150077}$) secondary antibody was used at 1/2000. Isotype control - Rabbit monoclonal lgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-GST3 / GST pi antibody [EPR8263] (ab138491)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of mouse stomach tissue sections labeling GST3 / GST pi with Purified ab138491 at 1/100 dilution (0.99 µg/mL). Heat mediated antigen retrieval using Bond™ Epitope Retrieval Solution 1 (pH 6.0). Rabbit specific IHC polymer detection kit HRP/DAB (ab209101) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-GST3 / GST pi antibody [EPR8263] (ab138491)


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human tonsil tissue sections labeling GST3 / GST pi with Purified ab138491 at 1/100 dilution (0.99 µg/mL). Heat mediated antigen retrieval using Bond™ Epitope Retrieval Solution 1 (pH 6.0). Rabbit specific IHC polymer detection kit HRP/DAB (ab209101) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.



Ol-RD Scanning - Anti-GST3 / GST pi antibody [EPR8263] (ab138491)

Equilibrium disassociation constant (K_D) Learn more about K_D

Click here to learn more about K_D



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors