

Anti-HCN2 antibody [S71] ab84817

★★★★★ **4 Abreviews** **10 References** 画像数 5

製品の概要

製品名	Anti-HCN2 antibody [S71]
製品の詳細	Mouse monoclonal [S71] to HCN2
由来種	Mouse
特異性	Does not cross react with HCN1.
アプリケーション	適用あり: IHC-Fr, IHC-P, ICC, WB
種交差性	交差種: Mouse, Rat, Human
免疫原	Fusion protein corresponding to Rat HCN2 aa 750 to the C-terminus (C terminal). Database link: Q9JKA9
ポジティブ・コントロール	IHC-Fr: Mouse brain tissue; IHC-P: Human hippocampus tissue; WB: Rat brain membrane and mouse brain tissue lysates; ICC: Differentiated SH-SH5Y cells.
特記事項	<p>The clone number S71 has previously been called S71-37 and N71/37. These clone numbers all refer to the same antibody clone.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid repeated freeze / thaw cycles.
バッファー	Preservative: 0.09% Sodium azide Constituents: 50% Glycerol, PBS
精製度	Protein G purified
ポリ/モノ	モノクローナル
クローン名	S71
アイソタイプ	IgG1

アプリケーション


The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab84817の使用に適用されます**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-Fr		1/1000.
IHC-P		1/100.
ICC		1/100.
WB	★★★★★ (3)	1/1000. Predicted molecular weight: 95 kDa.

ターゲット情報

機能	Hyperpolarization-activated ion channel exhibiting weak selectivity for potassium over sodium ions. Contributes to the native pacemaker currents in heart (If) and in neurons (Ih). Produces a large instantaneous current. Activated by cAMP. Modulated by intracellular chloride ions and pH; acidic pH shifts the activation to more negative voltages.
組織特異性	Highly expressed throughout the brain. Detected at low levels in heart.
配列類似性	Belongs to the potassium channel HCN family. Contains 1 cyclic nucleotide-binding domain.
ドメイン	The segment S4 is probably the voltage-sensor and is characterized by a series of positively charged amino acids at every third position.
細胞内局在	Membrane.

画像



Immunocytochemistry - Anti-HCN2 antibody [S71]
(ab84817)

Immunocytochemistry analysis of differentiated SH-SY5Y (human neuroblastoma cell line from bone marrow) cells labelling HCN2 with ab84817 at 1/100 dilution followed by secondary antibody (Alexa Fluor® 488) (green). Cells were counterstained with phalloidin (Alexa Fluor® 647) (red), and with Anti-beta III Tubulin antibody (Alexa Fluor® 555) (magenta). Nuclear DNA was labelled with Hoechst (blue).


(A) Phalloidin

(B) Anti-beta III Tubulin antibody

(C) HCN2 Antibody (ab84817) and secondary antibody


(D) Hoechst

(E) Composite




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HCN2 antibody [S71]
(ab84817)

Immunohistochemistry (Bouin-fixed paraffin-embedded sections) analysis of human hippocampus tissue labelling HCN2 with ab84817 at 1/100 dilution (1 hour at room temperature). FITC Goat Anti-Mouse (green) was used as the secondary antibody at 1/50 dilution (1 hour at room temperature).



Immunohistochemistry (Frozen sections) - Anti-HCN2 antibody [S71] (ab84817)

Immunohistochemistry (Frozen sections) analysis of mouse brain tissue labelling HCN2 with ab84817 at 1/1000 dilution (1 hour at room temperature). Tissue fixed with 10% Formalin Solution for 12-24 hours at room temperature. Secondary Antibody: HRP/DAB Detection System: Biotinylated Goat Anti-Mouse, Streptavidin Peroxidase, DAB Chromogen (brown) for 30 minutes at room temperature. Counterstained with Mayer hematoxylin (purple/blue) nuclear stain at 250-500 µL for 5 minutes at room temperature.



Anti-HCN2 antibody [S71] (ab84817) at 1/1000 dilution + Mouse brain tissue lysate at 15 µg


Secondary

Goat anti-mouse IgG, HRP at 1/4000 dilution

Predicted band size: 95 kDa

Observed band size: 95 kDa

Chemiluminescent for HRP (Moss) for 5 min in RT.



Anti-HCN2 antibody [S71] (ab84817) at 1/1000 dilution + Rat brain membrane lysates at 15 µg

Secondary

HRP-Sheep Anti-Mouse IgG antibody

Predicted band size: 95 kDa

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors