

Anti-HDAC2 antibody [Y461] ab32117

KO 評価済 リコンビナント RabMAb

★★★★★ **11 Abreviews** **83 References** 画像数 16

製品の概要

製品名	Anti-HDAC2 antibody [Y461]
製品の詳細	Rabbit monoclonal [Y461] to HDAC2
由来種	Rabbit
アプリケーション	適用あり: WB, IHC-P, ICC/IF, IP, Flow Cyt (Intra), ChIC/CUT&RUN-seq
種交差性	交差種: Mouse, Rat, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: HAP1, A431, Hela and K562 cell lysate and rat brain tissue homogenate. IHC-P: Human breast carcinoma and rat spinal cord tissue. ICC/IF: MCF-7 and wildtype HAP1 cells. Flow Cyt (intra): HeLa cells. ChIC/CUT&RUN-Seq: K-562 cells.
特記事項	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
バッファー	<p>pH: 7.20</p> <p>Preservative: 0.01% Sodium azide</p> <p>Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA</p>
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	Y461
アイソタイプ	IgG

アプリケーション


The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab32117の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB	★★★★★ (9)	1/2000. Predicted molecular weight: 55 kDa.
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol. Purified ab32117 at 1:1500 dilution.
ICC/IF		1/250 - 1/500.
IP		1/60.
Flow Cyt (Intra)		1/60 - 1/100. ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
ChIC/CUT&RUN-seq		Use at an assay dependent concentration. 5 µg

ターゲット情報

機能	Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes. Forms transcriptional repressor complexes by associating with MAD, SIN3, YY1 and N-COR. Interacts in the late S-phase of DNA-replication with DNMT1 in the other transcriptional repressor complex composed of DNMT1, DMAP1, PCNA, CAF1. Deacetylates TSHZ3 and regulates its transcriptional repressor activity.
組織特異性	Widely expressed; lower levels in brain and lung.
配列類似性	Belongs to the histone deacetylase family. HD type 1 subfamily.
翻訳後修飾	S-nitrosylated by GAPDH. In neurons, S-Nitrosylation at Cys-262 and Cys-274 does not affect the enzyme activity but abolishes chromatin-binding, leading to increases acetylation of histones and activate genes that are associated with neuronal development. In embryonic cortical neurons, S-Nitrosylation regulates dendritic growth and branching.
細胞内局在	Nucleus.

画像



Western blot - Anti-HDAC2 antibody [Y461]
(ab32117)

All lanes : Anti-HDAC2 antibody [Y461] (ab32117) at 1/1000 dilution

Lane 1 : HT-22 (mouse hippocampal neuronal cell) whole cell lysate

Lane 2 : SW10 (mouse neuronal Schwann cell) whole cell lysate

Lane 3 : bEnd.3 (mouse brain endothelioma) whole cell lysate

Lysates/proteins at 20 µg per lane.

Secondary


All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/100000 dilution

Predicted band size: 55 kDa

Observed band size: 60 kDa

Exposure time: 26 seconds

Blocking and diluting buffer and concentration: 5% NFDM/TBST.



Western blot - Anti-HDAC2 antibody [Y461]
(ab32117)

All lanes : Anti-HDAC2 antibody [Y461] (ab32117) at 1/1000 dilution

Lane 1 : GH3 (rat pituitary epithelial cell) whole cell lysate

Lane 2 : L6 (rat skeletal muscle myoblast) whole cell lysate

Lane 3 : C6 (rat glial tumor glial cell) whole cell lysate

Lane 4 : AR42J (rat pancreatic tumor epithelial cell) whole cell lysate

Lane 5 : 2.4G2 (rat B cell lymphoma B lymphocyte) whole cell lysate

Lysates/proteins at 20 µg per lane.

Secondary


All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/100000 dilution

Predicted band size: 55 kDa

Observed band size: 60 kDa

Exposure time: 37 seconds

Blocking and diluting buffer and concentration: 5% NFDM/TBST



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HDAC2 antibody [Y461] (ab32117)


Immunohistochemical analysis of paraffin-embedded fixed (A) Wild-type HEK293T (human embryonic kidney epithelial cell) cell pellet. (B) HDAC2 knockout HEK293T ([ab266590](#)) cell pellet staining HDAC2 with ab32117 at 1/10000 dilution followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Counter-staining used was hematoxylin.

Positive staining on (A) Wild-type HEK293T cell pellet, no staining on HDAC2 knockout HEK293T ([ab266590](#)) cell pellet.

The section was incubated with ab32117 for 30 mins at room temperature.

The immunostaining was performed on a Leica Biosystems BOND® RX instrument


Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins.



Immunocytochemistry/ Immunofluorescence - Anti-HDAC2 antibody [Y461] (ab32117)

Immunocytochemistry/ Immunofluorescence analysis of Wild-type HEK293T/HDAC2 KO HEK293T (HDAC2 knockout human embryonic kidney epithelial cell) ([ab266590](#)) cells labeling HDAC2 with ab32117 at 1/200 dilution followed by [ab150081](#) Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed secondary antibody at 1/1000 dilution (2 µg/ml). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Cells were counterstained with [ab195889](#) Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) at 1/200 dilution (2.5 µg/ml). DAPI was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.

Confocal image showing nuclear staining in Parental HEK293T cell line. Image was taken with a confocal microscope(Leica-Microsystems, TCS SP8).




ChIC/CUT&RUN sequencing - Anti-HDAC2 antibody [Y461] (ab32117)

ChIC/CUT&RUN was performed using a pAG-MNase at a final concentration of 700 ng/mL, 2×10^5 K-562 (Human chronic myelogenous leukemia lymphoblast) cells and 5 μ g of ab32117 [Y461]. The resulting DNA was sequenced on the Illumina NovaSeq 6000 to a depth of 10 million reads. The negative IgG control **ab172730** is also shown.

Additional screenshots of mapped reads can be downloaded [here](#).

The University of Geneva owns patents relevant to ChIC (Chromatin Immuno-Cleavage) methods.



Western blot - Anti-HDAC2 antibody [Y461] (ab32117)

All lanes : Anti-HDAC2 antibody [Y461] (ab32117) at 1/10000 dilution (Purified)

Lane 1 : HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate


Lane 2 : Mouse brain lysate

Lane 3 : Rat brain lysate

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/20000 dilution

Predicted band size: 55 kDa



Immunoprecipitation - Anti-HDAC2 antibody [Y461]
(ab32117)

Purified ab32117 at 1:20 dilution (0.5 µg) immunoprecipitating HDAC2 in HeLa whole cell lysate.

Lane 1 (input): HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate 10 µg.

Lane 2 (+): ab32117 + HeLa whole cell lysate.


Lane 3 (-): Rabbit monoclonal IgG (**ab172730**) instead of **ab32127** in HeLa whole cell lysate.

VeriBlot for IP Detection Reagent (HRP)(**ab131366**) (1:5000 dilution) was used for Western blotting.

Blocking Buffer and concentration: 5% NFDM/TBST.


Diluting buffer and concentration: 5% NFDM/TBST.

Observed band size: kDa




Flow Cytometry (Intracellular) - Anti-HDAC2 antibody
[Y461] (ab32117)

Flow Cytometry analysis of K-562 (Human chronic myelogenous leukemia lymphoblast) cells labelling HDAC2 with Purified ab32117 at 1:20 dilution (5 µg/ml) (Red). Cells were fixed with 4% Paraformaldehyde and permeabilised with 90% Methanol. A Goat anti rabbit IgG (Alexa Fluor® 488, **ab150081**) secondary antibody was used at 1:2000. Isotype control - Rabbit monoclonal IgG (Black). Unlabelled control - Cell without incubation with primary antibody and secondary antibody (Blue).




Immunocytochemistry/ Immunofluorescence - Anti-HDAC2 antibody [Y461] (ab32117)

Immunocytochemistry analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling HDAC2 with Purified ab32117 at 1:50 dilution (2.1 µg/ml). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Cells were counterstained with Ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1:200 (2.5 µg/ml). Goat anti rabbit IgG (Alexa Fluor® 488, **ab150077**) was used as the secondary antibody at 1:1000 (2 µg/ml) dilution. DAPI (blue) was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HDAC2 antibody [Y461] (ab32117)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of rat testis tissue sections labeling HDAC2 with Purified ab32117 at 1:1500 dilution (0.071 µg/ml). Heat mediated antigen retrieval was performed using Heat mediated antigen retrieval using Bond™ Epitope Retrieval Solution 2 (pH 9.0) . Tissue was counterstained with Hematoxylin. Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**) secondary antibody was used at 1:0 dilution. PBS instead of the primary antibody was used as the negative control.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HDAC2 antibody [Y461] (ab32117)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of mouse testis tissue sections labeling HDAC2 with Purified ab32117 at 1:1500 dilution (0.071 µg/ml). Heat mediated antigen retrieval was performed using Heat mediated antigen retrieval using Bond™ Epitope Retrieval Solution 2 (pH 9.0) . Tissue was counterstained with Hematoxylin. Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**) secondary antibody was used at 1:0 dilution. PBS instead of the primary antibody was used as the negative control.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HDAC2 antibody [Y461] (ab32117)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human testis tissue sections labeling HDAC2 with Purified ab32117 at 1:1500 dilution (0.071 µg/ml). Heat mediated antigen retrieval was performed using Heat mediated antigen retrieval using Bond™ Epitope Retrieval Solution 2 (pH 9.0) . Tissue was counterstained with Hematoxylin. Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**) secondary antibody was used at 1:0 dilution. PBS instead of the primary antibody was used as the negative control.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HDAC2 antibody [Y461] (ab32117)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human breast cancer tissue sections labeling HDAC2 with Purified ab32117 at 1:1500 dilution (0.071 µg/ml). Heat mediated antigen retrieval was performed using Heat mediated antigen retrieval using Bond™ Epitope Retrieval Solution 2 (pH 9.0) . Tissue was counterstained with Hematoxylin. Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**) secondary antibody was used at 1:0 dilution. PBS instead of the primary antibody was used as the negative control.



Western blot - Anti-HDAC2 antibody [Y461]
(ab32117)

Lane 1: Wild type HAP1 whole cell lysate (20 µg)


Lane 2: HDAC2 knockout HAP1 whole cell lysate (20 µg)

Lane 3: A431 whole cell lysate (20 µg)

Lane 4: HeLa whole cell lysate (20 µg)

Lanes 1 - 4: Merged signal (red and green). Green - ab32117 observed at 60 kDa. Red - loading control, **ab8245**, observed at 37 kDa.

ab32117 was shown to specifically react with HDAC2 when HDAC2 knockout samples were used. Wild-type and HDAC2 knockout samples were subjected to SDS-PAGE. ab32117 and **ab8245** (Mouse anti GAPDH loading control) were incubated overnight at 4°C at 1/2000 and 1/10000 respectively. Blots were developed with 800CW Goat anti Rabbit and 680CW Goat anti Mouse secondary antibodies at 1/10000 dilution for 1 hour at room temperature before imaging.



Immunocytochemistry/ Immunofluorescence - Anti-HDAC2 antibody [Y461] (ab32117)

ab32117 staining HDAC2 in wild-type HAP1 cells (top panel) and HDAC2 knockout HAP1 cells (bottom panel). The cells were fixed with 100% methanol (5min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated with ab32117 at 1/250 dilution and **ab195889** at 1/250 dilution (shown in pseudocolour red) overnight at +4°C, followed by a further incubation at room temperature for 1h with a goat secondary antibody to Rabbit IgG (Alexa Fluor® 488) (**ab150081**) at 2 µg/ml (shown in green). Nuclear DNA was labelled in blue with DAPI.

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-HDAC2 antibody [Y461] (ab32117)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors