


### Anti-Histone H1.0 antibody [34] ab11079

KO 評価済

★★★★☆ 2 Abreviews 31 References 画像数 6

#### 製品の概要

製品名	Anti-Histone H1.0 antibody [34]
製品の詳細	Mouse monoclonal [34] to Histone H1.0
由来種	Mouse
アプリケーション	適用あり: WB, IHC-P
種交差性	交差種: Human, Recombinant fragment 交差が予測される動物種: Mouse, Rat, Xenopus laevis, Vertebrata  非交差種: Bird
免疫原	Full length native protein (purified) corresponding to Cow Histone H1.0.
エピトープ	This antibody recognises an epitope within aa20-30. Proline 26 is not essential for recognition by the antibody. See Gorka et al. 1998 for more information.
ポジティブ・コントロール	WB: A431, MCF7 and HeLa cell lysate; Histone H1.0 (Human) - Recombinant Protein. IHC-P: Human breast adenocarcinoma tissue.
特記事項	<p>This antibody clone is manufactured by Abcam. If you require a custom buffer formulation or conjugation for your experiments, please contact <a href="mailto:orders@abcam.com">orders@abcam.com</a>.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

#### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	pH: 7.40 Preservative: 0.02% Sodium azide Constituents: PBS, 6.97% L-Arginine

精製度	Protein G purified
ポリ/モノ	モノクローナル
クローン名	34
ミエローマ	NS1/1-Ag4-1
アイソタイプ	IgG1

アプリケーション


**The Abpromise guarantee**      **Abpromise保証**は、次のテスト済みアプリケーションにおけるab11079の使用に適用されます  
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB	★★★★★ (2)	Use a concentration of 1 µg/ml. Detects a band of approximately 32 kDa (predicted molecular weight: 20 kDa).
IHC-P		Use a concentration of 1 µg/ml. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.

ターゲット情報

機能	Histones H1 are necessary for the condensation of nucleosome chains into higher-order structures. The H1F0 histones are found in cells that are in terminal stages of differentiation or that have low rates of cell division.
配列類似性	Belongs to the histone H1/H5 family. Contains 1 H15 (linker histone H1/H5 globular) domain.
翻訳後修飾	Phosphorylated on Ser-17 in RNA edited version.
細胞内局在	Nucleus. Chromosome. The RNA edited version has been localized to nuclear speckles. During mitosis, it appears in the vicinity of condensed chromosomes.

画像



Western blot - Anti-Histone H1.0 antibody [34] (ab11079)

**All lanes :** Anti-Histone H1.0 antibody [34] (ab11079) at 1/500 dilution

**Lane 1 :** Wild-type A431 cell lysate

**Lane 2 :** H1F0 knockout A431 cell lysate

**Lane 3 :** MCF7 (Human breast adenocarcinoma cell line) whole cell lysate

Lysates/proteins at 40 µg per lane.


Performed under reducing conditions.

**Predicted band size:** 20 kDa

**Observed band size:** 30 kDa

**Lanes 1 - 3:** Merged signal (red and green). Green - ab11079 observed at 30 kDa. Red - loading control **ab52866** (Rabbit anti-alpha Tubulin antibody [EP1332Y]) observed at 55 kDa.

ab11079 was shown to react with Histone H1.0 in wild-type A431 cells in Western blot with loss of signal observed in H1F0 knockout sample. Wild-type A431 and H1F0 knockout cell lysates were subjected to SDS-PAGE. Membranes were blocked in 3 % milk in TBS-T (0.1 % Tween®) before incubation with ab11079 and **ab52866** (Rabbit anti-alpha Tubulin antibody [EP1332Y]) overnight at 4°C at a 1 in 500 dilution and a 1 in 20000 dilution respectively. Blots were incubated with Goat anti-Mouse IgG H&L (IRDye® 800CW) preabsorbed (**ab216772**) and Goat anti-Rabbit IgG H&L (IRDye® 680RD) preabsorbed (**ab216777**) secondary antibodies at 1 in 20000 dilution for 1 h at room temperature before imaging.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Histone H1.0 antibody [34] (ab11079)

IHC image of Histone H1.0 staining in a formalin fixed, paraffin embedded human breast adenocarcinoma tissue section\*, performed on a Leica Bond™ system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab11079, 5µg/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.

\*Tissue obtained from the Human Research Tissue Bank, supported by the NIHR Cambridge Biomedical Research Centre



Western blot - Anti-Histone H1.0 antibody [34] (ab11079)

**All lanes :** Anti-Histone H1.0 antibody [34] (ab11079) at 1 µg/ml

**Lane 1 :** Histone H1.0 (Human) - Recombinant Protein

**Lane 2 :** Histone H1.2 (Human) - Recombinant Protein

Lysates/proteins at 0.1 µg per lane.

### Secondary


**All lanes :** Goat polyclonal to Mouse IgG - H&L - Pre-Adsorbed (HRP) ([ab65485](#)) at 1/5000 dilution

Performed under reducing conditions.

**Predicted band size:** 20 kDa

**Observed band size:** 32 kDa

**Exposure time:** 10 seconds



Western blot - Anti-Histone H1.0 antibody [34]  
(ab11079)

Anti-Histone H1.0 antibody [34] (ab11079) at 1 µg/ml + HeLa  
(Triton-enriched) Cell Lysate at 10 µg

### Secondary

Goat polyclonal to Mouse IgG - H&L - Pre-Adsorbed (HRP)  
([ab65485](#)) at 1/5000 dilution

Performed under reducing conditions.

**Predicted band size:** 20 kDa

**Observed band size:** 32 kDa

**Additional bands at:** 120 kDa, 18 kDa. We are unsure as to the  
identity of these extra bands.


**Exposure time:** 8 minutes

Western blot - Anti-Histone H1.0 antibody [34]  
(ab11079)

Western blot using ab11079 on HeLa whole cell extract (30  
µg/lane).

Exposure 5 sec using ECL.

12.5% SDS-PAGE gel.



Mouse monoclonal to Histone H1.0 against recombinant human H1 isoforms H1.0 to H1.5 as depicted in the figure.

Dilution 1/1000.

Recombinant H1 isoforms courtesy of Dr Nicole Happel.

(see attached review)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors