

# Anti-Histone H4 antibody [mAbcam 17036] - ChIP Grade ab17036

★★★★★ [4 Abreviews](#) [9 References](#) [画像数 7](#)

### 製品の概要

製品名	Anti-Histone H4 antibody [mAbcam 17036] - ChIP Grade
製品の詳細	Mouse monoclonal [mAbcam 17036] to Histone H4 - ChIP Grade
由来種	Mouse
アプリケーション	<b>適用あり:</b> IHC-P, ChIP, WB
種交差性	<b>交差種:</b> Mouse, Rat, Cow, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers. (Peptide available as <a href="#">ab13843</a> )
ポジティブ・コントロール	This antibody gave a positive signal in the following: Calf thymus histone preparation; HeLa histone preparation; Histone H4 recombinant protein; NIH3T3 Whole Cell Lysate; MEF1 Whole Cell Lysate; PC12 Whole Cell Lysate.
特記事項	<p>To detect histone H4 by ICC/IF, we recommend using <a href="#">ab177840</a> or <a href="#">ab9052</a> since these antibodies perform better in this application.</p> <p>This antibody clone is manufactured by Abcam. If you require a custom buffer formulation or conjugation for your experiments, please contact <a href="mailto:orders@abcam.com">orders@abcam.com</a>.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	pH: 7.40 Preservative: 0.02% Sodium azide

	Constituent: PBS
精製度	IgG fraction
ポリ/モノ	モノクローナル
クローン名	mAbcam 17036
ミエローマ	Sp2/0-Ag14
アイソタイプ	IgG2a
軽鎖の種類	kappa

アプリケーション

The Abpromise guarantee      Abpromise保証は、次のテスト済みアプリケーションにおけるab17036の使用に適用されます  
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-P	★★★★★ (1)	Use a concentration of 1 µg/ml.
ChIP		Use 5 µg for 25 µg of chromatin.
WB	★★★★★ (2)	Use a concentration of 1 µg/ml. Detects a band of approximately 14 kDa (predicted molecular weight: 11 kDa).Can be blocked with <b>Human Histone H4 peptide (ab13843)</b> . We recommend using 3% milk as the blocking agent for Western blot.

ターゲット情報

機能	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.
配列類似性	Belongs to the histone H4 family.
翻訳後修飾	Acetylation at Lys-6 (H4K5ac), Lys-9 (H4K8ac), Lys-13 (H4K12ac) and Lys-17 (H4K16ac) occurs in coding regions of the genome but not in heterochromatin. Citruination at Arg-4 (H4R3ci) by PAD4 impairs methylation. Monomethylation and asymmetric dimethylation at Arg-4 (H4R3me1 and H4R3me2a, respectively) by PRMT1 favors acetylation at Lys-9 (H4K8ac) and Lys-13 (H4K12ac). Demethylation is performed by JMJD6. Symmetric dimethylation on Arg-4 (H4R3me2s) by the PRDM1/PRMT5 complex may play a crucial role in the germ-cell lineage. Monomethylated, dimethylated or trimethylated at Lys-21 (H4K20me1, H4K20me2, H4K20me3). Monomethylation is performed by SET8. Trimethylation is performed by SUV420H1 and SUV420H2 and induces gene silencing. Ubiquitinated by the CUL4-DDB-RBX1 complex in response to ultraviolet irradiation. This may weaken the interaction between histones and DNA and facilitate DNA accessibility to repair proteins. Monoubiquitinated at Lys-92 of histone H4 (H4K91ub1) in response to DNA damage. The exact role of H4K91ub1 in DNA damage response is still unclear but it may function as a


licensing signal for additional histone H4 post-translational modifications such as H4 Lys-21 methylation (H4K20me).

Sumoylated, which is associated with transcriptional repression.


細胞内局在

Nucleus. Chromosome.

## 画像




Chromatin was prepared from HeLa cells according to the Abcam X-ChIP protocol. Cells were fixed with formaldehyde for 10 minutes. The ChIP was performed with 25µg of chromatin, 5µg of ab17036 (blue), and 20µl of protein A/G sepharose beads. No antibody was added to the beads control (yellow). The immunoprecipitated DNA was quantified by real time PCR (Taqman approach for active and inactive loci, Sybr green approach for heterochromatic loci). Primers and probes are located in the first kb of the transcribed region.



IHC image of ab17036 staining in Human Hodgekin's Lymphoma formalin fixed paraffin embedded tissue section, performed on a Leica Bond™ system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab17036, 1µg/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.



**All lanes :** Anti-Histone H4 antibody [mAbcam 17036] - ChIP Grade (ab17036) at 2 µg/ml

**Lane 1 :** Histone H1 Recombinant Protein

**Lane 2 :** Histone H2A Recombinant Protein

**Lane 3 :** Histone H2B Recombinant Protein

**Lane 4 :** Histone H3.1 Recombinant Protein

**Lane 5 :** Histone H4 Recombinant Protein

Lysates/proteins at 0.1 µg per lane.

### Secondary


**All lanes :** Goat polyclonal to Mouse IgG - H&L - Pre-Adsorbed (HRP) at 1/5000 dilution

Performed under reducing conditions.

**Predicted band size:** 11 kDa

**Observed band size:** 11 kDa

**Exposure time:** 30 seconds



**All lanes :** Anti-Histone H4 antibody [mAbcam 17036] - ChIP Grade (ab17036) at 1 µg/ml

**Lane 1 :** NIH 3T3 (Mouse embryonic fibroblast cell line) Whole Cell Lysate

**Lane 2 :** MEF1 (Mouse embryonic fibroblast cell line) Whole Cell Lysate

**Lane 3 :** PC12 (Rat adrenal pheochromocytoma cell line) Whole Cell Lysate

Lysates/proteins at 10 µg per lane.

### Secondary

**All lanes :** Goat Anti-Mouse IgG H&L (HRP) preadsorbed (**ab97040**) at 1/5000 dilution


Developed using the ECL technique.

Performed under reducing conditions.

**Predicted band size:** 11 kDa

**Observed band size:** 14 kDa

**Exposure time:** 1 minute



Western blot - Anti-Histone H4 antibody [mAbcam 17036] - ChIP Grade (ab17036)

**All lanes :** Anti-Histone H4 antibody [mAbcam 17036] - ChIP

Grade (ab17036) at 5 µg/ml

**Lane 1 :** Calf Thymus Histone Preparation Nuclear Lysate at 0.5 µg

**Lane 2 :** HeLa Histone Preparation Nuclear Lysate at 2.5 µg

#### Secondary


**All lanes :** Goat polyclonal to Mouse IgG - H&L - Pre-Adsorbed (HRP) at 1/3000 dilution

Performed under reducing conditions.

**Predicted band size:** 11 kDa

**Observed band size:** 14 kDa

**Exposure time:** 1 minute




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Histone H4 antibody [mAbcam 17036] - ChIP Grade (ab17036)

IHC image of Histone H4 staining in formalin fixed, paraffin embedded human Hodgkin's lymphoma tissue section\*, performed on a Leica Bond™ system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab17036, 1 µg/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.

\*Tissue obtained from the Human Research Tissue Bank,  
supported by the NIHR Cambridge Biomedical Research Centre



**All lanes :** Anti-Histone H4 antibody [mAbcam 17036] - ChIP  
Grade (ab17036) at 5 µg/ml

**Lane 1 :** Histone H1 Recombinant Protein

**Lane 2 :** Histone H2A Recombinant Protein

**Lane 3 :** Histone H2B Recombinant Protein

**Lane 4 :** Histone H3.1 Recombinant Protein

**Lane 5 :** Histone H4 Recombinant Protein

Lysates/proteins at 0.1 µg per lane.

### Secondary

**All lanes :** Goat polyclonal to Mouse IgG - H&L - Pre-Adsorbed  
(HRP) at 1/3000 dilution

Performed under reducing conditions.

**Predicted band size:** 11 kDa

**Observed band size:** 14 kDa

**Exposure time:** 30 seconds

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors