

Anti-HLA G antibody [MEM-G/1] ab7759

★★★★★ **3 Abreviews** **25 References** 画像数 4

製品の概要

製品名	Anti-HLA G antibody [MEM-G/1]
製品の詳細	Mouse monoclonal [MEM-G/1] to HLA G
由来種	Mouse
特異性	The antibody reacts with denaturated HLA-G heavy chain.
アプリケーション	適用あり: IHC-P, WB
種交差性	交差種: Human
免疫原	Recombinant full length denatured heavy chain (Human).
ポジティブ・コントロール	IHC-P: First trimester placenta and pulmonary diseases. WB: ab29745 ; HeLa and JEG-3 cell lysates.
特記事項	<p>This product was changed from ascites to tissue culture supernatant on 24th January 2018. Please note that the dilutions may need to be adjusted accordingly. If you have any questions, please do not hesitate to contact our scientific support team.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	<p>pH: 7.40</p> <p>Preservative: 0.097% Sodium azide</p> <p>Constituent: PBS</p>
精製度	Protein A purified
特記事項 (精製)	Purified from TCS. Purity >95% by SDS-PAGE.

ポリ/モノ	モノクローナル
クローン名	MEM-G/1
アイソタイプ	IgG1

アプリケーション


The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab7759の使用に適用されます
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-P	★★★★★ (1)	1/60 - 1/100. Perform heat mediated antigen retrieval via the microwave method before commencing with IHC staining protocol.
WB	★★★★★ (1)	Use a concentration of 1 - 10 µg/ml. Predicted molecular weight: 38.2 kDa.

ターゲット情報


機能	Involved in the presentation of foreign antigens to the immune system. Plays a role in maternal tolerance of the fetus by mediating protection from the deleterious effects of natural killer cells, cytotoxic T lymphocytes, macrophages and mononuclear cells.
組織特異性	Expressed in trophoblasts.
配列類似性	Belongs to the MHC class I family. Contains 1 Ig-like C1-type (immunoglobulin-like) domain.
細胞内局在	Membrane.

画像



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HLA G antibody [MEM-G/1] (ab7759)

Immunohistochemistry staining with ab7759 of first trimester placenta (paraffin-embedded sections).



Western blot - Anti-HLA G antibody [MEM-G/1] (ab7759)

All lanes : Anti-HLA G antibody [MEM-G/1] (ab7759) at 10 µg/ml

Lane 1 : Human placenta tissue lysate - total protein ([ab29745](#))

Lane 2 : HeLa (Human epithelial carcinoma cell line) Whole Cell Lysate

Lane 3 : JEG-3 (Human placental choriocarcinoma cell line) Whole Cell Lysate

Lysates/proteins at 10 µg per lane.


Secondary

All lanes : Goat polyclonal to Mouse IgG - H&L - Pre-Adsorbed (HRP) at 1/3000 dilution

Predicted band size: 38.2 kDa


Observed band size: 42 kDa

Additional bands at: 38 kDa. We are unsure as to the identity of these extra bands.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HLA G antibody [MEM-G/1] (ab7759)

The antibody ab7759 stains infiltrating macrophages in pulmonary diseases. In the top left corner see detail of macrophage.



Western blot - Anti-HLA G antibody [MEM-G/1] (ab7759)

All lanes : Anti-HLA G antibody [MEM-G/1] (ab7759) at 2 µg/ml

Lane 1 : JEG-3 cell line under reducing conditions.

Lane 2 : LNCaP cell line under reducing conditions.

Lane 3 : JEG-3 cell line under non-reducing conditions.

Lane 4 : LNCaP cell line under non-reducing conditions.

Secondary

All lanes : IRDye800-conjugated anti-mouse secondary antibody

Predicted band size: 38.2 kDa

Observed band size: 40 kDa

Western blotting analysis of human HLA-G using ab7759 on lysates of JEG-3 cell line and LNCaP cell line (negative control) under reducing and non-reducing conditions.

Nitrocellulose membrane was probed with 2 µg/ml of ab7759 followed by IRDye800-conjugated anti-mouse secondary antibody.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you

- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors