

HRP Anti-Caspase-7 antibody [E22] ab206039

KO 評価済 RabMAb

画像数 2

製品の概要

製品名	HRP Anti-Caspase-7 antibody [E22]
製品の詳細	HRP Rabbit monoclonal [E22] to Caspase-7
由来種	Rabbit
標識	HRP
アプリケーション	適用あり: WB
種交差性	交差種: Human
免疫原	Synthetic peptide within Human Caspase-7 aa 1-100 (N terminal). The exact sequence is proprietary. Database link: P55210
ポジティブ・コントロール	WB: Jurkat whole cell lysate.
特記事項	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C.

	Avoid freeze / thaw cycle. Store In the Dark.
バッファー	pH: 7.40 Preservative: 0.1% Proclin 300 Solution Constituents: 30% Glycerol (glycerin, glycerine), 1% BSA, PBS
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	E22
アイソタイプ	IgG

アプリケーション


The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab206039の使用に適用されます**
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		1/5000. Detects a band of approximately 36 kDa (predicted molecular weight: 34 kDa).

ターゲット情報

機能	Involved in the activation cascade of caspases responsible for apoptosis execution. Cleaves and activates sterol regulatory element binding proteins (SREBPs). Proteolytically cleaves poly(ADP-ribose) polymerase (PARP) at a '216-Asp-Gly-217' bond. Overexpression promotes programmed cell death.
組織特異性	Highly expressed in lung, skeletal muscle, liver, kidney, spleen and heart, and moderately in testis. No expression in the brain.
配列類似性	Belongs to the peptidase C14A family.
翻訳後修飾	Cleavages by granzyme B or caspase-10 generate the two active subunits. Propeptide domains can also be cleaved efficiently by caspase-3. Active heterodimers between the small subunit of caspase-7 and the large subunit of caspase-3, and vice versa, also occur.
細胞内局在	Cytoplasm.

画像



All lanes : HRP Anti-Caspase-7 antibody [E22] (ab206039) at 1/5000 dilution

Lane 1 : Wild-type HAP1 whole cell lysate

Lane 2 : CASP7 knockout HAP1 whole cell lysate

Lysates/proteins at 20 µg per lane.

Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 34 kDa

Observed band size: 38 kDa

Exposure time: 20 minutes

ab206039 was shown to specifically react with Caspase-7 in wild-type HAP1 cells as signal was lost in CASP7 knockout cells. Wild-type and CASP7 knockout samples were subjected to SDS-PAGE. Ab206039 was incubated overnight at 4°C at 1/5000 dilution. Blots were developed with ECL technique.



Western blot - HRP Anti-Caspase-7 antibody [E22]
(ab206039)

HRP Anti-Caspase-7 antibody [E22] (ab206039) at 1/5000 dilution
+ Jurkat (Human T cell lymphoblast-like cell line) Whole Cell Lysate
at 10 µg

Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 34 kDa

Observed band size: 36 kDa

Exposure time: 2 minutes

This blot was produced using a 4-12% Bis-tris gel under the MOPS buffer system.

The gel was run at 200V for 50 minutes before being transferred onto a Nitrocellulose membrane at 30V for 70 minutes. The membrane was then blocked for an hour using 2% Bovine Serum Albumin before being incubated with ab206039 overnight at 4°C.

Antibody binding was visualised using ECL development solution [ab133406](#).

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors