# abcam

### Product datasheet

## HRP Anti-elF4E antibody [Y448] ab205824

יעלאעבע RabMAb

#### 画像数 2

#### 製品の概要

製品名 HRP Anti-elF4E antibody [Y448]

製品の詳細 HRP Rabbit monoclonal [Y448] to elF4E

由来種 Rabbit 標識 HRP

アプリケーション **適用あり:** WB 種交差性 交差種: Human

交差が予測される動物種: Mouse, Rat 4

免疫原 Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

ポジティブ・コントロール WB: HEK293 whole cell lysate.

特記事項 This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity

- Long-term security of supply

- Animal-free production

For more information **see here**.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

#### 製品の特性

製品の状態 Liquid

Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. 保存方法

Stable for 12 months at -20°C. Store In the Dark.

バッファー pH: 7.40

Preservative: 0.1% Proclin 300 Solution

Constituents: 30% Glycerol (glycerin, glycerine), 1% BSA, PBS

精製度 Protein A purified

ポリモノ モノクローナル

クローン名 Y448

#### アプリケーション


**The Abpromise guarantee** <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab205824の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		1/5000. Detects a band of approximately 30 kDa (predicted molecular weight: 25 kDa).

#### ターゲット情報

機能	Its translation stimulation activity is repressed by binding to the complex CYFIP1-FMR1 (By similarity). Recognizes and binds the 7-methylguanosine-containing mRNA cap during an early step in the initiation of protein synthesis and facilitates ribosome binding by inducing the unwinding of the mRNAs secondary structures. Component of the CYFIP1-EIF4E-FMR1 complex which binds to the mRNA cap and mediates translational repression. In the CYFIP1-EIF4E-FMR1 complex this subunit mediates the binding to the mRNA cap.
配列類似性	Belongs to the eukaryotic initiation factor 4E family.
翻訳後修飾	Phosphorylation increases the ability of the protein to bind to mRNA caps and to form the elF4F complex.

#### 画像



HRP Anti-elF4E antibody [Y448] (ab205824) at 1/5000 dilution + HEK293 (Human embryonic kidney cell line) Whole Cell Lysate at 10  $\mu g$ 

Developed using the ECL technique.


Performed under reducing conditions.

**Predicted band size:** 25 kDa **Observed band size:** 30 kDa

Exposure time: 12 minutes

This blot was produced using a 4-12% Bis-tris gel under the MES buffer system. The gel was run at 200V for 35 minutes before being transferred onto a Nitrocellulose membrane at 30V for 70 minutes.

The membrane was then blocked for an hour using 3% milk before being incubated with ab205824 overnight at 4°C. Antibody binding was visualised using ECL development solution <u>ab133406</u>.



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- · Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.co.jp/abpromise">https://www.abcam.co.jp/abpromise</a> or contact our technical team.

#### Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors