abcam

Product datasheet

HRP Anti-IL-13 antibody [EP3266] ab201470

ועלשעבע RabMAb

1 References 画像数 2

製品の概要

製品名 HRP Anti-IL-13 antibody [EP3266]

製品の詳細 HRP Rabbit monoclonal [EP3266] to IL-13

由来種 Rabbit HRP 標識

アプリケーション **適用あり:** WB 種交差性 交差種: Human

交差が予測される動物種: Mouse, Rat 4

免疫原 Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

ポジティブ・コントロール WB: Human IL-13 recombinant protein.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit 特記事項

monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C.

Avoid freeze / thaw cycle. Store In the Dark.

バッファー pH: 7.40

Preservative: 0.1% Proclin 300 Solution

Constituents: 30% Glycerol (glycerin, glycerine), PBS, 1% BSA

精製度 Protein A purified

ポリモノ モノクローナル

クローン名 EP3266

アイソタイプ ΙgG

アプリケーション

Abpromise保証は、次のテスト済みアプリケーションにおけるab201470の使用に適用されます The Abpromise guarantee

アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

Secreted.

アプリケーション	Abreviews	特記事項
WB		1/5000. Detects a band of approximately 13 kDa (predicted molecular weight: 13 kDa).


ターゲット情報

機能
Cytokine (PubMed:8096327, PubMed:8097324). Inhibits inflammatory cytokine production (PubMed:8096327). Synergizes with IL2 in regulating interferon-gamma synthesis (PubMed:8096327). May be critical in regulating inflammatory and immune responses (PubMed:8096327, PubMed:8097324). Positively regulates IL31RA expression in macrophages.

関連疾患
配列類似性
Belongs to the IL-4/IL-13 family.

画像

細胞内局在



Western blot - HRP Anti-IL-13 antibody [EP3266] (ab201470)

HRP Anti-IL-13 antibody [EP3266] (ab201470) at 1/5000 dilution + Human IL-13 Recombinant Protein at 0.1 µg


Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 13 kDa **Observed band size:** 13 kDa

Exposure time: 2 seconds

This blot was produced using a 4-12% Bis-tris gel under the MES buffer system. The gel was run at 200V for 35 minutes before being transferred onto a Nitrocellulose membrane at 30V for 70 minutes. The membrane was then blocked for an hour using 2% Bovine Serum Albumin before being incubated with ab201470 overnight at 4°C. Antibody binding was visualised using ECL development solution ab133406.



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors