abcam

Product datasheet

HRP Anti-Metabotropic Glutamate Receptor 5 antibody [EPR2425Y] ab196482

יעלטעבעו RabMAb

画像数 2

製品の概要

製品名 HRP Anti-Metabotropic Glutamate Receptor 5 antibody [EPR2425Y]

製品の詳細 HRP Rabbit monoclonal [EPR2425Y] to Metabotropic Glutamate Receptor 5

由来種 Rabbit 標識 HRP

アプリケーション 適用あり: IHC-P 種交差性 交差種: Human

交差が予測される動物種: Mouse, Rat 🔷

免疫原 Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

ポジティブ・コントロール IHC-P: Normal human cerebellum tissue.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit 特記事項

monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C.

Avoid freeze / thaw cycle. Store In the Dark.

バッファー pH: 7.40

Preservative: 0.1% Proclin 300 Solution

Constituents: 30% Glycerol (glycerin, glycerine), 1% BSA, PBS

精製度 Protein A purified

ポリモノ モノクローナル クローン名 EPR2425Y

アイソタイプ ΙgG

アプリケーション

The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab196482の使用に適用されます


アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-P		1/100. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. ab199507 - Rabbit monoclonal IgG (HRP), is suitable for use an as isotype control with this antibody.

ターゲット情報

機能	G-protein coupled receptor for glutamate. Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of down-stream effectors. Signaling activates a phosphatidylinositol-calcium second messenger system and generates a calcium-activated chloride current. Plays an important role in the regulation of synaptic plasticity and the modulation of the neural network activity.
配列類似性	Belongs to the G-protein coupled receptor 3 family.
細胞内局在	Cell membrane.

画像



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - HRP Anti-Metabotropic
Glutamate Receptor 5 antibody [EPR2425Y]
(ab196482)

IHC image of Metabotropic Glutamate Receptor 5 staining in a section of formalin-fixed paraffin-embedded normal human cerebellum, performed on a Leica BOND. The section was pretreated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20mins. The section was then incubated with ab196482 at 1/100 dilution, for 15 mins at room temperature. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX. The inset negative control image is taken from an identical assay without primary antibody.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors