abcam

Product datasheet

Anti-HSV1 ICP0 antibody [5H7] ab6513

**** ★ 1 Abreviews 39 References 画像数 1

製品の概要

種交差性

製品名 Anti-HSV1 ICP0 antibody [5H7]

製品の詳細 Mouse monoclonal [5H7] to HSV1 ICP0

由来種 Mouse

アプリケーション 適用あり: WB, ELISA, ICC/IF

免疫原 Herpes Virus

特記事項

The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

交差種: Herpes simplex virus

If you have any questions, special requirements or concerns, please send us an inquiry and/or

contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw

cycles.

バッファー pH: 7.4

精製度 Protein A purified

ポリ/モノ モノクローナル

クローン名 5H7

₹**I**□-マ NS1/1-Ag4-1

アイソタイプ lgG1

軽鎖の種類 kappa

アプリケーション

The Abpromise guarantee

Abpromise保証は、次のテスト済みアプリケーションにおけるab6513の使用に適用されます

アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		1/8000.
ELISA		1/12800.
ICC/IF	**** <u>(1)</u>	1/25600.

ターゲット情報


関連性

Herpes simplex virus (HSV) encodes four nuclear phosphoproteins that contribute towards the full productive lytic cycle cascade in newly infected permissive cells. The IE175 (or ICP4) and IE63 (or ICP27) proteins have essential regulatory roles during the lytic cycle. A third, IE110 (or ICP0), appears to play only an accessory role in the presence of VP16, but in the absence of VP16 appears to be critical for initiation of reactivated infections.

細胞内局在

Nuclear

画像



Immunocytochemistry/ Immunofluorescence - Anti-HSV1 ICP0 antibody [5H7] (ab6513)

This image is courtesy of an anonymous Abreview

ab6513 (1/25600) staining HSV1 ICP0 in human cells. Cells were PFA-fixed, permeabilized in 0.3% TritonX-100/PBS and blocked in 1% serum /30 minutes /25°C. A fluorescently labeled donkey antimouse antibody was used as the secondary.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery

- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors