

Anti-IKK alpha antibody [EPR464] ab109749

KO 評価済 リコンビナント RabMAb

7 References 画像数 5

製品の概要

製品名	Anti-IKK alpha antibody [EPR464]
製品の詳細	Rabbit monoclonal [EPR464] to IKK alpha
由来種	Rabbit
アプリケーション	適用あり: WB, IP, IHC-P 適用なし: Flow Cyt (Intra) or ICC/IF
種交差性	交差種: Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: Jurkat, HeLa, HepG2 and Daudi cell lysates; IHC: Human breast tissue IP: Jurkat cell lysate.
特記事項	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p> <p>Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.
バッファー	pH: 7.20 Preservative: 0.05% Sodium azide Constituents: 0.1% BSA, 40% Glycerol (glycerin, glycerine), 9.85% Tris glycine, 50% Tissue culture supernatant
精製度	Protein A purified
ポリ/モノ	モノクローナル

クローン名	EPR464
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee **Abpromise保証は、** 次のテスト済みアプリケーションにおけるab109749の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。


アプリケーション	Abreviews	特記事項
WB		1/1000 - 1/10000. Predicted molecular weight: 85 kDa.
IP		1/10 - 1/100.
IHC-P		1/100 - 1/250. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

追加情報 Is unsuitable for Flow Cyt (Intra) or ICC/IF.

ターゲット情報

機能	Acts as part of the IKK complex in the conventional pathway of NF-kappa-B activation and phosphorylates inhibitors of NF-kappa-B thus leading to the dissociation of the inhibitor/NF-kappa-B complex and ultimately the degradation of the inhibitor. As part of the non-canonical pathway of NF-kappa-B activation, the MAP3K14-activated CHUK/IKKA homodimer phosphorylates NFkB2/p100 associated with RelB, inducing its proteolytic processing to NFkB2/p52 and the formation of NF-kappa-B RelB-p52 complexes. Also phosphorylates NCOA3. Phosphorylates 'Ser-10' of histone H3 at NF-kappa-B-regulated promoters during inflammatory responses triggered by cytokines.
組織特異性	Widely expressed.
関連疾患	Defects in CHUK are the cause of cocoon syndrome (COCOS) [MIM:613630]; also known as fetal encasement syndrome. COCOS is a lethal syndrome characterized by multiple fetal malformations including defective face and seemingly absent limbs, which are bound to the trunk and encased under the skin.
配列類似性	Belongs to the protein kinase superfamily. Ser/Thr protein kinase family. I-kappa-B kinase subfamily. Contains 1 protein kinase domain.
翻訳後修飾	Phosphorylated by MAP3K14/NIK, AKT and to a lesser extent by MEKK1, and dephosphorylated by PP2A. Autophosphorylated. Acetylation of Thr-179 by Yersinia yopJ prevents phosphorylation and activation, thus blocking the I-kappa-B signaling pathway.
細胞内局在	Cytoplasm. Nucleus. Shuttles between the cytoplasm and the nucleus.

画像



Western blot - Anti-IKK alpha antibody [EPR464]
(ab109749)

Lanes 1, 3 and 5: Wild-type HAP1 cell lysate (20 µg)


Lanes 2, 4 and 6: IKK alpha knockout HAP1 cell lysate (20 µg)

Lanes 1 and 2: Green signal from target - ab109749 observed at 84 kDa

Lanes 3 and 4: Red signal from loading control - **ab8226** observed at 42 kDa

Lanes 5 and 6: Merged (red and green) signal

ab109749 was shown to specifically react with IKK alpha when IKK alpha knockout samples were used. Wild-type and IKK alpha knockout samples were subjected to SDS-PAGE. ab109749 and **ab8226** (loading control to beta actin) were both diluted at 1/1000 and incubated overnight at 4°C. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (**ab216776**) secondary antibodies at 1/10000 dilution for 1 h at room temperature before imaging.



Western blot - Anti-IKK alpha antibody [EPR464]
(ab109749)

All lanes : Anti-IKK alpha antibody [EPR464] (ab109749) at 1/1000 dilution

Lane 1 : Jurkat cell lysate


Lane 2 : HeLa cell lysate

Lane 3 : HepG2 cell lysate

Lane 4 : Daudi cell lysate

Lysates/proteins at 10 µg per lane.

Predicted band size: 85 kDa



Immunoprecipitation - Anti-IKK alpha antibody
[EPR464] (ab109749)


IKK alpha was immunoprecipitated from 0.35 mg Jurkat (Human T cell leukemia T lymphocyte) whole cell lysate 10 µg with 109749 at 1/50 dilution (2µg). VeriBlot for IP Detection Reagent (HRP) ([ab131366](#)) was used at 1/5000 dilution.

Lane 1: Jurkat (Human T cell leukemia T lymphocyte) whole cell lysate 10 µg

Lane 2: ab109749 IP in Jurkat whole cell lysate

Lane 3: Rabbit monoclonal IgG ([ab172730](#)) instead of ab109749 in Jurkat whole cell lysate

Blocking and dilution buffer and concentration: 5% NFDm/TBST.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-IKK alpha antibody
[EPR464] (ab109749)

ab109749, at 1/100 dilution, staining IKK alpha in paraffin-embedded Human breast tissue by Immunohistochemistry.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-IKK alpha antibody [EPR464] (ab109749)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors