

Anti-IL-10RA antibody [1B1.3a] ab33738

1 References

製品の概要

製品名	Anti-IL-10RA antibody [1B1.3a]
製品の詳細	Rat monoclonal [1B1.3a] to IL-10RA
由来種	Rat
アプリケーション	適用あり: Flow Cyt
種交差性	交差種: Mouse 非交差種: Human
免疫原	Recombinant fragment corresponding to Mouse IL-10RA.
エピトープ	The epitope recognised by this antibody has been localised to the ligand binding domain within the extracellular region of mouse CD210 (IL-10RA).
特記事項	<p>This clone is reported to neutralize the effects of IL-10.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term.
バッファー	pH: 7.40 Preservative: 0.09% Sodium azide Constituent: PBS
精製度	Protein G purified
一次抗体 備考	This clone is reported to neutralize the effects of IL10.
ポリ/モノ	モノクローナル

クローン名	1B1.3a
ミエローマ	Sp2/0
アイソタイプ	IgG1

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab33738の使用に適用されます**
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt		Use at an assay dependent concentration. Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul. ab18407 - Rat monoclonal IgG1, is suitable for use as an isotype control with this antibody.

ターゲット情報

機能	Receptor for IL10; binds IL10 with a high affinity.
組織特異性	Spleen, thymus, and PBMC. Weak expression in pancreas, skeletal muscle, brain, heart, and kidney. Placenta, lung, and liver showed intermediate levels. Monocytes, B-cells, large granular lymphocytes, and T-cells express high levels.
関連疾患	Inflammatory bowel disease 28
配列類似性	Belongs to the type II cytokine receptor family.
細胞内局在	Membrane.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors