

Anti-IL-17A antibody ab9565

★★★★★ **2 Abreviews** **19 References**

製品の概要

製品名	Anti-IL-17A antibody
製品の詳細	Rabbit polyclonal to IL-17A
由来種	Rabbit
アプリケーション	適用あり: IHC-P, ELISA, WB
種交差性	交差種: Human
免疫原	Recombinant full length protein corresponding to Human IL-17A. Highly pure (>98%) recombinant hIL-17 (human Interleukin-17) Database link: Q16552
ポジティブ・コントロール	WB: Recombinant human IL-17A protein (ab9567).
特記事項	<p>Although some customers have been successful in IHC we no longer batch test in this application.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Lyophilized:Reconstitute with 200µl of sterile water. Please note that if you receive this product in liquid form it has already been reconstituted as described and no further reconstitution is necessary.
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term.
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	unknown
軽鎖の種類	unknown

アプリケーション

The Abpromise guarantee Abpromise保証は、 次のテスト済みアプリケーションにおけるab9565の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-P	★★★★★ (2)	Use at an assay dependent concentration. PubMed: 29390966
ELISA		Use at an assay dependent concentration. Allows the detection of 0.2 - 0.4 ng/well of recombinant hIL-17 (assuming 100 µl/well antibody solution).
WB		Use at an assay dependent concentration.

ターゲット情報

機能	Induces stromal cells to produce proinflammatory and hematopoietic cytokines. Enhances the surface expression of the intracellular adhesion molecule-1 (ICAM-1) in fibroblasts.
組織特異性	Restricted to activated memory T-cells.
配列類似性	Belongs to the IL-17 family.
翻訳後修飾	Found both in glycosylated and nonglycosylated forms.
細胞内局在	Secreted.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors