


### Anti-IL-33 antibody ab118503

[5 References](#) [画像数 4](#)

#### 製品の概要

製品名	Anti-IL-33 antibody
製品の詳細	Rabbit polyclonal to IL-33
由来種	Rabbit
アプリケーション	<b>適用あり:</b> WB, IHC-P, ELISA
種交差性	<b>交差種:</b> Human <b>交差が予測される動物種:</b> Mouse, Rat 
免疫原	Synthetic peptide corresponding to Human IL-33 (internal sequence) conjugated to keyhole limpet haemocyanin.
ポジティブ・コントロール	Human lymph node lysate and uterus tissue.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

#### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term.
バッファー	pH: 7.4 Preservative: 0.02% Sodium azide Constituent: 99% PBS
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

#### アプリケーション

## The Abpromise guarantee

**Abpromise保証は、次のテスト済みアプリケーションにおけるab118503の使用に適用されます**


アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		Use a concentration of 1 µg/ml. Predicted molecular weight: 31 kDa.
IHC-P		Use a concentration of 5 µg/ml. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
ELISA		Use at an assay dependent concentration.

## ターゲット情報


機能	<p>Cytokine that binds to and signals through the IL1RL1/ST2 receptor which in turn activates NF-kappa-B and MAPK signaling pathways in target cells (PubMed:16286016). Involved in the maturation of Th2 cells inducing the secretion of T-helper type 2-associated cytokines. Also involved in activation of mast cells, basophils, eosinophils and natural killer cells. Acts as a chemoattractant for Th2 cells, and may function as an "alarmin", that amplifies immune responses during tissue injury (PubMed:17853410, PubMed:18836528).</p> <p>In quiescent endothelia the uncleaved form is constitutively and abundantly expressed, and acts as a chromatin-associated nuclear factor with transcriptional repressor properties, it may sequester nuclear NF-kappaB/RELA, lowering expression of its targets (PubMed:21734074). This form is rapidly lost upon angiogenic or proinflammatory activation (PubMed:18787100).</p>
組織特異性	Expressed at high level in high endothelial venules found in tonsils, Peyer patches and mesenteric lymph nodes. Almost undetectable in placenta.
配列類似性	Belongs to the IL-1 family. Highly divergent.
ドメイン	The homeodomain-like HTH domain mediates nuclear localization and heterochromatin association.
翻訳後修飾	The full length protein can be released from cells and is able to signal via the IL1RL1/ST2 receptor. However, proteolytic processing by CSTG/cathepsin G and ELANE/neutrophil elastase produces C-terminal peptides that are more active than the unprocessed full length protein. May also be proteolytically processed by calpains (PubMed:19596270). Proteolytic cleavage mediated by apoptotic caspases including CASP3 and CASP7 results in IL33 inactivation (PubMed:19559631). In vitro proteolytic cleavage by CASP1 was reported (PubMed:16286016) but could not be confirmed in vivo (PubMed:19465481) suggesting that IL33 is probably not a direct substrate for that caspase.
細胞内局在	Nucleus. Chromosome. Cytoplasmic vesicle, secretory vesicle. Secreted. Associates with heterochromatin and mitotic chromosomes (PubMed:17185418).

## 画像




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-IL-33 antibody (ab118503)

Immunohistochemical staining of paraffin-embedded, formalin fixed, human tonsil lymphocytes with ab118503 at 5 µg/mL.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-IL-33 antibody (ab118503)

Negative control - immunohistochemical staining of paraffin-embedded, formalin fixed, human seminiferous tubules with ab118503 at 5 µg/mL.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-IL-33 antibody (ab118503)

Human Uterus, Vessels (formalin-fixed, paraffin-embedded) stained with IL33 antibody ab118503 at 5 µg/ml followed by biotinylated goat anti-rabbit IgG secondary antibody, alkaline phosphatase-streptavidin and chromogen.



**Lane 1** : Anti-IL-33 antibody (ab118503) at 1 µg/ml

**Lane 2** : Anti-IL-33 antibody (ab118503) at 2 µg/ml

**All lanes** : Human lymph node tissue lysate

**Predicted band size:** 31 kDa

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors