

# Anti-Insulin antibody [EPR17359] ab181547

リコンビナント **RabMAb**

★★★★★ **11 Abreviews** **76 References** 画像数 **11**

### 製品の概要

製品名	Anti-Insulin antibody [EPR17359]
製品の詳細	Rabbit monoclonal [EPR17359] to Insulin
由来種	Rabbit
特異性	The human recommendation is based on the IHC-P results. We do not guarantee WB for human.
アプリケーション	<b>適用あり:</b> IHC-FoFr, WB, IHC-P, ICC/IF, mlHC
種交差性	<b>交差種:</b> Mouse, Rat, Human
免疫原	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	IHC-P: Human, mouse and rat pancreas tissue. ICC/IF: BxPC-3 cells. IHC-FoFr: Mouse pancreas tissue. WB: Mouse & Rat pancreas lysate. mlHC: Human pancreas tissue.
特記事項	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> <li>- Animal-free production</li> </ul> <p>For more information <a href="#">see here</a>.</p> <p>Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb<sup>®</sup> patents</a>.</p>

### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol, 0.05% BSA
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EPR17359
アイソタイプ	IgG

## アプリケーション


**The Abpromise guarantee** **Abpromise保証は、** 次のテスト済みアプリケーションにおけるab181547の使用に適用されます  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-FoFr		1/1000.
WB	★★★★★ (1)	Use at an assay dependent concentration. Predicted molecular weight: 12 kDa.
IHC-P	★★★★★ (7)	1/64000. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
ICC/IF		1/200.
mlHC		1/20000.

## ターゲット情報

機能	Insulin decreases blood glucose concentration. It increases cell permeability to monosaccharides, amino acids and fatty acids. It accelerates glycolysis, the pentose phosphate cycle, and glycogen synthesis in liver.
関連疾患	<p>Defects in INS are the cause of familial hyperproinsulinemia (FHPRI) [MIM:176730].</p> <p>Defects in INS are a cause of diabetes mellitus insulin-dependent type 2 (IDDM2) [MIM:125852]. IDDM2 is a multifactorial disorder of glucose homeostasis that is characterized by susceptibility to ketoacidosis in the absence of insulin therapy. Clinical features are polydipsia, polyphagia and polyuria which result from hyperglycemia-induced osmotic diuresis and secondary thirst. These derangements result in long-term complications that affect the eyes, kidneys, nerves, and blood vessels.</p> <p>Defects in INS are a cause of diabetes mellitus permanent neonatal (PNDM) [MIM:606176]. PNDM is a rare form of diabetes distinct from childhood-onset autoimmune diabetes mellitus type 1. It is characterized by insulin-requiring hyperglycemia that is diagnosed within the first months of life. Permanent neonatal diabetes requires lifelong therapy.</p> <p>Defects in INS are a cause of maturity-onset diabetes of the young type 10 (MODY10) [MIM:613370]. MODY10 is a form of diabetes that is characterized by an autosomal dominant mode of inheritance, onset in childhood or early adulthood (usually before 25 years of age), a primary defect in insulin secretion and frequent insulin-independence at the beginning of the disease.</p>
配列類似性	Belongs to the insulin family.
細胞内局在	Secreted.

## 画像




Fluorescence multiplex immunohistochemical analysis of the human pancreas (Formalin/PFA-fixed paraffin-embedded sections).

Panel A: merged staining of anti-Carboxypeptidase A ([ab278044](#), magenta; Opal™690), anti-Cytokeratin 19 ([ab195872](#), green; Opal™520) and anti-Insulin (ab181547, red; Opal™570) on human pancreas. Panel B: anti-Carboxypeptidase A stained on acinar cells. Panel C: anti-Cytokeratin 19 stained on centroacinar cells and ducts. Panel D: anti-Insulin stained on beta cells. Opal Polymer HRP Ms + Rb was used as a secondary antibody.

The section was incubated in three rounds of staining: in the order of [ab278044](#) at 1/4000 dilution (0.135 µg/ml), [ab195872](#) at 1/8000 dilution (0.127 µg/ml), and ab181547 at 1/20000 dilution (0.053 µg/ml) for 30 mins at room temperature. Each round was followed by a separate fluorescent tyramide signal amplification system.

The immunostaining was performed on a Leica Biosystems BOND® RX instrument with an Opal™ 4-color kit. Image acquisition was performed with Leica SP8 confocal microscope.

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins. DAPI (blue) was used as a nuclear counter stain.



**All lanes :** Anti-Insulin antibody [EPR17359] (ab181547) at 1/1000 dilution

**Lane 1 :** Human pancreas

**Lane 2 :** Mouse pancreas

**Lane 3 :** Rat pancreas

Lysates/proteins at 20 µg per lane.


### Secondary

**All lanes :** Goat Anti-Rabbit IgG (HRP) with minimal cross-reactivity with human IgG at 2000 µg/ml

**Predicted band size:** 12 kDa

Blocking/Diluting buffer and concentration: 5% NFDM/TBST

Observed MW: 12KDa



Western blot - Anti-Insulin antibody [EPR17359]  
(ab181547)


Anti-Insulin antibody [EPR17359] (ab181547) at 1/1000 dilution +  
Mouse pancreas lysate 20 µg at 20 µg

### Secondary

Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

**Predicted band size:** 12 kDa

**Observed band size:** 12 kDa




Immunohistochemistry (Formalin/PFA-fixed paraffin-  
embedded sections) - Anti-Insulin antibody  
[EPR17359] (ab181547)

Immunohistochemical analysis of paraffin-embedded human  
pancreas tissue labeling Insulin with ab181547 at 1/64000 dilution,  
followed by Anti-Rabbit HRP ([ab97051](#)) at 1/500 dilution.

Cytoplasm staining on islet cells of human pancreas is  
observed. Counterstained with hematoxylin.

**Negative control:** PBS instead of primary antibody; secondary  
antibody is Anti-Rabbit HRP ([ab97051](#)) at 1/500 dilution.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH  
9.0 before commencing with IHC staining protocol.



Immunocytochemistry/ Immunofluorescence - Anti-Insulin antibody [EPR17359] (ab181547)


Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized BxPC-3 (Human pancreas adenocarcinoma cells) cells labeling Insulin with ab181547 at 1/200 dilution, followed by Goat anti-rabbit IgG (Alexa Fluor® 488) (**ab150077**) secondary antibody at 1/400 dilution (green).

Confocal image shows cytoplasmic staining on BxPC-3 cells. The nuclear counterstain is DAPI (blue). Tubulin is detected with **ab7291** (anti-Tubulin mouse mAb) at 1/500 dilution and **ab150120** (Alexa Fluor® 594 Goat anti-Mouse secondary) at 1/500 dilution (red).

The negative controls are as follows:-

**-ve control 1:** - ab181547 at 1/200 dilution followed by **ab150120** (Alexa Fluor® 594 Goat anti-Mouse secondary) at 1/500 dilution.

**-ve control 2:** - **ab7291** (anti-Tubulin mouse mAb) at 1/500 dilution followed by **ab150077** (Alexa Fluor® 488 Goat Anti-Rabbit IgG H&L) at 1/400 dilution.




Immunohistochemistry (PFA perfusion fixed frozen sections) - Anti-Insulin antibody [EPR17359] (ab181547)

Immunohistochemical analysis of 4% paraformaldehyde perfusion fixed, frozen section of mouse pancreas tissue labeling Insulin with ab181547 at 1/1000 dilution, followed by Donkey anti-rabbit Alexa Fluor® 594 at 1/1000 dilution. Cytoplasm staining on islet cells of mouse pancreas is observed. Counter stained with DAPI.

**Negative control:** PBS instead of primary antibody; secondary antibody is Donkey anti-rabbit Alexa Fluor® 594 at 1/1000 dilution.






Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Insulin antibody [EPR17359] (ab181547)

Immunohistochemical analysis of paraffin-embedded mouse pancreas tissue labeling Insulin with ab181547 at 1/64000 dilution, followed by Anti-Rabbit HRP ([ab97051](#)) at 1/500 dilution. Cytoplasm staining on islet cells of mouse pancreas is observed. Counterstained with hematoxylin.

**Negative control:** PBS instead of primary antibody; secondary antibody is Anti-Rabbit HRP ([ab97051](#)) at 1/500 dilution.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Insulin antibody [EPR17359] (ab181547)

Immunohistochemical analysis of paraffin-embedded rat pancreas tissue labeling Insulin with ab181547 at 1/64000 dilution, followed by Anti-Rabbit HRP ([ab97051](#)) at 1/500 dilution. Cytoplasm staining on islet cells of rat pancreas is observed. Counterstained with hematoxylin.

**Negative control:** PBS instead of primary antibody; secondary antibody is Anti-Rabbit HRP ([ab97051](#)) at 1/500 dilution.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Insulin antibody [EPR17359] (ab181547)

Immunohistochemical analysis of paraffin-embedded human adenocarcinoma of colon tissue with ab181547 at 1/64000 dilution, followed by Anti-Rabbit HRP (**ab97051**) at 1/500 dilution.

**Negative staining** on human colonic adenocarcinoma is observed. Counterstained with hematoxylin.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Insulin antibody [EPR17359] (ab181547)

Immunohistochemical analysis of paraffin-embedded human liver tissue with ab181547 at 1/64000 dilution, followed by Anti-Rabbit HRP (**ab97051**) at 1/500 dilution.

**Negative staining** on human liver tissue is observed. Counterstained with hematoxylin.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

### Why choose a recombinant antibody?



Anti-Insulin antibody [EPR17359] (ab181547)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors