

Anti-Integrin alpha 4/CD49D (phospho S988) antibody [EPR2343] ab75760

リコンビナント RabMAb

★★★★☆ **1 Abreviews** **1 References** **画像数 4**

製品の概要

製品名	Anti-Integrin alpha 4/CD49D (phospho S988) antibody [EPR2343]
製品の詳細	Rabbit monoclonal [EPR2343] to Integrin alpha 4/CD49D (phospho S988)
由来種	Rabbit
特異性	ab75760 only detects Integrin alpha 4/CD49D phosphorylated on Serine 988 of the mature protein. S988 corresponds to S1021 of the sequence with the 33 aa signal peptide and S1027 of the precursor shown in Figure 4 of PMID: 8436405. The antibody detects the mature chain (180 kDa), the precursor (140 kDa) and the cleaved C terminal (70 kDa).

アプリケーション

適用あり: WB
適用なし: ICC/IF or IHC-P

種交差性

交差種: Mouse, Rat, Human

免疫原

Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

ポジティブ・コントロール

THP1 cell lysates

特記事項

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information [see here](#).

Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to [RabMAb[®] patents](#).

製品の特性

製品の状態

Liquid

保存方法

Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Stable for 12 months at -20°C.

バッファー	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EPR2343
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab75760の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。


アプリケーション	Abreviews	特記事項
WB	★★★★★ (1)	1/500 - 1/2000. Predicted molecular weight: 150 kDa.

追加情報 Is unsuitable for ICC/IF or IHC-P.

ターゲット情報

機能	Integrins alpha-4/beta-1 (VLA-4) and alpha-4/beta-7 are receptors for fibronectin. They recognize one or more domains within the alternatively spliced CS-1 and CS-5 regions of fibronectin. They are also receptors for VCAM1. Integrin alpha-4/beta-1 recognizes the sequence Q-I-D-S in VCAM1. Integrin alpha-4/beta-7 is also a receptor for MADCAM1. It recognizes the sequence L-D-T in MADCAM1. On activated endothelial cells integrin VLA-4 triggers homotypic aggregation for most VLA-4-positive leukocyte cell lines. It may also participate in cytolytic T-cell interactions with target cells.
配列類似性	Belongs to the integrin alpha chain family. Contains 7 FG-GAP repeats.
ドメイン	The SG1 motif is involved in binding to chondroitin sulfate glycosaminoglycan and cell adhesion.
翻訳後修飾	Phosphorylation on Ser-1027 inhibits PXN binding.
細胞内局在	Membrane.

画像



Western blot - Anti-Integrin alpha 4/CD49D (phospho S988) antibody [EPR2343] (ab75760)

All lanes : Anti-Integrin alpha 4/CD49D (phospho S988) antibody [EPR2343] (ab75760) at 1/1000 dilution (purified)

Lane 1 : Rat spleen tissue lysate

Lane 2 : Mouse spleen tissue lysate

Lysates/proteins at 10 µg per lane.

Secondary


All lanes : HRP goat anti-rabbit IgG (H+L) at 1/1000 dilution

Predicted band size: 150 kDa

Observed band size: 150,70 kDa

Blocking buffer: 5% NFDm/TBST

Dilution buffer: 5% NFDm/TBST



Western blot - Anti-Integrin alpha 4/CD49D (phospho S988) antibody [EPR2343] (ab75760)

All lanes : Anti-Integrin alpha 4/CD49D (phospho S988) antibody [EPR2343] (ab75760) at 1/5000 dilution (purified)

Lane 1 : untreated Jurkat lysate

Lane 2 : Jurkat cell lysate treated with Lambda phosphatase

Lysates/proteins at 10 µg per lane.

Secondary


All lanes : HRP goat anti-rabbit IgG (H+L) at 1/50000 dilution

Predicted band size: 150 kDa

Observed band size: 140,150,70 kDa

Blocking buffer: 5% NFDm/TBST

Dilution buffer: 5% NFD/MTBST



Western blot - Anti-Integrin alpha 4/CD49D (phospho S988) antibody [EPR2343] (ab75760)

All lanes : Anti-Integrin alpha 4/CD49D (phospho S988) antibody [EPR2343] (ab75760) at 1/2000 dilution

Lane 1 : THP1 cell lysates, untreated

Lane 2 : THP1 cell lysates, with Lambda Phosphatase

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : goat anti-rabbit HRP at 1/2000 dilution

Predicted band size: 150 kDa

Observed band size: 140,180,70 kDa

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-Integrin alpha 4/CD49D (phospho S988) antibody [EPR2343] (ab75760)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors