# abcam

#### Product datasheet

## Anti-Integrin linked ILK antibody [EP1593Y] ab52480



ייבעדיו RabMAb

#### 18 References 画像数 10

#### 製品の概要

製品名 Anti-Integrin linked ILK antibody [EP1593Y]

製品の詳細 Rabbit monoclonal [EP1593Y] to Integrin linked ILK

由来種 Rabbit

特異性 This antibody reacts with Integrin linked ILK. The mouse and rat recommendation is based on the

WB results. We do not guarantee IHC-P for mouse and rat.

アプリケーション 適用あり: WB, IP, IHC-P

適用なし: Flow Cyt or ICC/IF

種交差性 交差種: Rat, Human, African green monkey

交差が予測される動物種: Mouse, Cow 🔷

免疫原 Synthetic peptide within Human Integrin linked ILK aa 400 to the C-terminus. The exact sequence

is proprietary.

Database link: Q13418

ポジティブ・コントロール Human heart muscle tissue and COS cell lysate

特記事項 This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity

- Long-term security of supply

- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

#### 製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long

term. Avoid freeze / thaw cycle.

バッファー pH: 7.20

Preservative: 0.01% Sodium azide

Constituents: 59% PBS, 40% Glycerol, 0.05% BSA

精製度 Protein A purified

**ポリ/モノ** モノクローナル

**クローン名** EP1593Y

アイソタイプ IgG

#### アプリケーション

The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab52480の使用に適用されますアプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		1/1000. Detects a band of approximately 51 kDa (predicted molecular weight: 51 kDa).  For unpurified use at 1/2000 - 1/10000 dilution.
IP		1/40. For unpurified use at 1/30 dilution.
IHC-P		1/100. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.  See IHC antigen retrieval protocols.  The mouse and rat recommendation is based on the WB results.  We do not guarantee IHC-P for mouse and rat.

追加情報 Is unsuitable for Flow Cyt or ICC/IF.

#### ターゲット情報

機能	Receptor-proximal protein kinase regulating integrin-mediated signal transduction. May act as a mediator of inside-out integrin signaling. Focal adhesion protein part of the complex ILK-PINCH. This complex is considered to be one of the convergence points of integrin- and growth factor-signaling pathway. Could be implicated in mediating cell architecture, adhesion to integrin substrates and anchorage-dependent growth in epithelial cells. Phosphorylates beta-1 and beta-3 integrin subunit on serine and threonine residues, but also AKT1 and GSK3B.
組織特異性	Highly expressed in heart followed by skeletal muscle, pancreas and kidney. Weakly expressed in placenta, lung and liver.

**配列類似性** Belongs to the protein kinase superfamily. TKL Ser/Thr protein kinase family.

Contains 5 ANK repeats.


Contains 1 protein kinase domain.

ドメイン A PH-like domain is involved in phosphatidylinositol phosphate binding.


**翻訳後修飾** Autophosphorylated on serine residues.

**細胞内局在** Cell junction > focal adhesion. Cell membrane.

### 画像



Western blot - Anti-Integrin linked ILK antibody [EP1593Y] (ab52480)



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Integrin linked ILK antibody [EP1593Y] (ab52480)

Lane 1: Wild-type HAP1 cell lysate (20 µg)

Lane 2: Integrin linked ILK knockout HAP1 cell lysate (20 µg)

Lane 3: HEK293 cell lysate (20 µg)

Lane 4: K562 cell lysate (20 µg)

**Lanes 1 - 4:** Merged signal (red and green). Green - ab52480 observed at 52 kDa. Red - loading control, <u>ab8245</u>, observed at 37 kDa.

Unpurified ab52480 was shown to specifically react with Integrin linked ILK when Integrin linked ILK knockout samples were used. Wild-type and Integrin linked ILK knockout samples were subjected to SDS-PAGE. Unpurified ab52480 and <a href="mailto:ab8245">ab8245</a> (loading control to GAPDH) were both diluted 1/2000 and incubated overnight at 4°C. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (<a href="mailto:ab216773">ab216773</a>) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (<a href="mailto:ab216776">ab216776</a>) secondary antibodies at 1/10000 dilution for 1 hour at room temperature before imaging.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human lung cancer tissue sections labeling Integrin linked ILK with Purified ab52480 at 1/100 dilution (7.13 µg/ml). Heat mediated antigen retrieval was performed using citrate (pH 6.0).



Western blot - Anti-Integrin linked ILK antibody [EP1593Y] (ab52480)

**All lanes :** Anti-Integrin linked ILK antibody [EP1593Y] (ab52480) at 1/1000 dilution (Purified)

**Lane 1**: HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates with 5% NFDM/TBST

**Lane 2 :** K-562 (Human chronic myelogenous leukemia lymphoblast) whole cell lysates with 5% NFDM/TBST

Lane 3: Mouse heart lysates with 5% NFDM/TBST

**Lane 4**: HEK-293 (Human embryonic kidney epithelial cell) whole cell lysates with 5% NFDM/TBST

**Lane 5**: PC-12 (Rat adrenal gland pheochromocytoma) whole cell lysates with 5% NFDM/TBST

Lysates/proteins at 20 µg per lane.

#### **Secondary**

**All lanes :** Goat Anti-Rabbit lgG H&L (HRP) (<u>ab97051</u>) at 1/20000 dilution (Goat Anti-Rabbit lgG, (H+L), Peroxidase conjugated)

**Predicted band size:** 51 kDa **Observed band size:** 51 kDa



Immunoprecipitation - Anti-Integrin linked ILK antibody [EP1593Y] (ab52480)

ab52480 (purified) at 1:40 dilution (2µg) immunoprecipitating Integrin linked ILK in HeLa whole cell lysate.

Lane 1 (input): HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate 10µg


Lane 2 (+): ab52480 & HeLa whole cell lysate

**Lane 3 (-):** Rabbit monoclonal IgG (<u>ab172730</u>) instead of ab52480 in HeLa whole cell lysate

For western blotting, VeriBlot for IP Detection Reagent (HRP)

(ab131366) was used for detection at 1:1000 dilution.

Blocking and diluting buffer: 5% NFDM/TBST.




Western blot - Anti-Integrin linked ILK antibody [EP1593Y] (ab52480)

Anti-Integrin linked ILK antibody [EP1593Y] (ab52480) at 1/1000 dilution (Purified) + COS-1 (African green monkey kidney fibroblast-like) whole cell lysates at 15 µg with 5% NFDM/TBST

#### **Secondary**

Goat Anti-Rabbit lgG H&L (HRP) (ab97051) at 1/20000 dilution (Goat Anti-Rabbit lgG, (H+L), Peroxidase conjugated)

**Predicted band size:** 51 kDa **Observed band size:** 51 kDa




Western blot - Anti-Integrin linked ILK antibody [EP1593Y] (ab52480) Anti-Integrin linked ILK antibody [EP1593Y] (ab52480) at 1/2000 dilution (unpurified) + COS-1 (African green monkey kidney fibroblast-like) cell lysate at 10 µg

#### Secondary

Goat anti-rabbit HRP labeled at 1/2000 dilution


**Predicted band size:** 51 kDa **Observed band size:** 51 kDa



Western blot - Anti-Integrin linked ILK antibody [EP1593Y] (ab52480)


Anti-Integrin linked ILK antibody [EP1593Y] (ab52480) (unpurified) + PC12

Predicted band size: 51 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Integrin linked ILK antibody [EP1593Y] (ab52480)

Immunohistochemical analysis of paraffin-embedded human heart muscle using unpurified ab52480 at a 1/50 dilution.




Western blot - Anti-Integrin linked ILK antibody [EP1593Y] (ab52480)

 $\begin{tabular}{ll} \textbf{All lanes:} Anti-Integrin linked ILK antibody [EP1593Y] (ab52480) \\ at 1/2000 dilution (unpurified) \\ \end{tabular}$ 

Lane 1: Rat Brain
Lane 2: Rat Heart
Lane 3: Rat Kidney
Lane 4: Rat Spleen

Predicted band size: 51 kDa

Exposure time: 3 minutes



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- · Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.co.jp/abpromise">https://www.abcam.co.jp/abpromise</a> or contact our technical team.

#### Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors