

# Anti-Interferon alpha/beta receptor 1 (phospho S535 + S539) antibody ab30571

画像数 1

### 製品の概要

製品名	Anti-Interferon alpha/beta receptor 1 (phospho S535 + S539) antibody
製品の詳細	Rabbit polyclonal to Interferon alpha/beta receptor 1 (phospho S535 + S539)
由来種	Rabbit
特異性	Specific for Interferon alpha/beta receptor 1 protein phosphorylated at Ser535,539 in Western blots of human brain extracts.
アプリケーション	<b>適用あり:</b> WB
種交差性	<b>交差種:</b> Human
免疫原	Synthetic peptide corresponding to Human Interferon alpha/beta receptor 1 (phospho S535 + S539).
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
バッファー	pH: 7.50 Constituents: 0.238% HEPES, 50% Glycerol, 0.87% Sodium chloride, 0.01% BSA
精製度	Immunogen affinity purified
特記事項 (精製)	Prepared from rabbit serum by affinity purification via sequential chromatography on phospho-and dephosphopeptide affinity columns.
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

## アプリケーション


**The Abpromise guarantee**      **Abpromise保証は、次のテスト済みアプリケーションにおけるab30571の使用に適用されます**  
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
WB		1/1000. Predicted molecular weight: 110-130 kDa. Note: the molecular weight of the IFNNAR1 varies with cell line (different levels of glycosylation) in 293 and HeLa Cells; the mature form is ~110 - 130k.

## ターゲット情報

機能	Associates with IFNAR2 to form the type I interferon receptor. Receptor for interferons alpha and beta. Binding to type I IFNs triggers tyrosine phosphorylation of a number of proteins including JAKs, TYK2, STAT proteins and IFNR alpha- and beta-subunits themselves.
組織特異性	IFN receptors are present in all tissues and even on the surface of most IFN-resistant cells. Isoform 1, isoform 2 and isoform 3 are expressed in the IFN-alpha sensitive myeloma cell line U266S. Isoform 2 and isoform 3 are expressed in the IFN-alpha resistant myeloma cell line U266R. Isoform 1 is not expressed in IFN-alpha resistant myeloma cell line U266R.
配列類似性	Belongs to the type II cytokine receptor family. Contains 3 fibronectin type-III domains.
翻訳後修飾	Phosphorylated on tyrosine residues by TYK2 tyrosine kinase.
細胞内局在	Membrane.

## 画像



**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors