

Anti-Kallikrein 2/KLK2 antibody [6B7] ab40749

1 References [画像数 1](#)

製品の概要

製品名	Anti-Kallikrein 2/KLK2 antibody [6B7]
製品の詳細	Mouse monoclonal [6B7] to Kallikrein 2/KLK2
由来種	Mouse
特異性	This antibody reacts with human Kallikrein 2/KLK2. This is no detectable reactivity with PSA.
アプリケーション	適用あり: IHC-P
種交差性	交差種: Human
免疫原	Tissue, cells or virus. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	IHC-P: Human prostate tissue.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term.
バッファー	Preservative: 0.1% Sodium azide Constituent: PBS
精製度	Protein G purified
ポリ/モノ	モノクローナル
クローン名	6B7
アイソタイプ	IgG1

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab40749の使用に適用されます**
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-P		

追加情報

AP: Use at an assay dependent dilution.
 ELISA: Use at an assay dependent dilution.
 ICC: Use at an assay dependent dilution.
 IHC-P: Use at an assay dependent dilution. As a guideline use at 18ug/ml for sections. Perform heat mediated antigen retrieval via the microwave method before commencing with IHC staining protocol.

Not yet tested in other applications.
 Optimal dilutions/concentrations should be determined by the end user.

ターゲット情報

機能 Glandular kallikreins cleave Met-Lys and Arg-Ser bonds in kininogen to release Lys-bradykinin.
配列類似性 Belongs to the peptidase S1 family. Kallikrein subfamily.
 Contains 1 peptidase S1 domain.

画像



ab40749 staining Kallikrein on Human Prostate Tissue by Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) at 18 µg/ml.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish

- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors