

Anti-Kazrin antibody ab88752

画像数 2

製品の概要

製品名	Anti-Kazrin antibody
製品の詳細	Mouse polyclonal to Kazrin
由来種	Mouse
アプリケーション	適用あり: WB
種交差性	交差種: Rat, Human
免疫原	Recombinant full length protein within Human Kazrin aa 1-450. The exact immunogen sequence used to generate this antibody is proprietary information. If additional detail on the immunogen is needed to determine the suitability of the antibody for your needs, please contact our Scientific Support team to discuss your requirements.
ポジティブ・コントロール	Rat brain tissue lysate and transfected 293T cell lysate
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
バッファー	pH: 7.40 Constituent: 100% PBS
精製度	Protein A purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee

Abpromise保証は、次のテスト済みアプリケーションにおけるab88752の使用に適用されず


アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		Use a concentration of 1 µg/ml. Predicted molecular weight: 86 kDa.

ターゲット情報

機能	Component of the cornified envelope of keratinocytes. May be involved in the interplay between adherens junctions and desmosomes. The function in the nucleus is not known.
組織特異性	Isoform 2, isoform 3 and isoform 4 are expressed in several cell lines including keratinocytes and bladder and epidermoid carcinoma (at protein level). Isoform 2, isoform 3 and isoform 4 are expressed in hair follicle and interfollicular epidermis (at protein level).
配列類似性	Belongs to the kazrin family. Contains 3 SAM (sterile alpha motif) domains.
細胞内局在	Cytoplasm > cytoskeleton and Cytoplasm. Cell junction > desmosome. Nucleus. Observed at the apical plasma mebrane of keratinocytes. Partially colocalizes with PPL and DP at desmosomes, and with PP at the interdesmosomal plasma membrane. Colocalizes with cortical actin-based membrane structures.

画像




Anti-Kazrin antibody (ab88752) at 1 µg/ml + Rat brain tissue lysate at 50ug/lane

Predicted band size: 86 kDa

Observed band size: 86 kDa

Additional bands at: 45 kDa. We are unsure as to the identity of these extra bands.



Western blot - Anti-Kazrin antibody (ab88752)

All lanes : Anti-Kazrin antibody (ab88752) at 1 µg/ml

Lane 1 : Kazrin transfected 293T cell lysate (45.65 kDa) at 25ug/lane

Lane 2 : Non-transfected 293T cell lysate at 25ug/lane

Predicted band size: 86 kDa

Observed band size: 60 kDa

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors