


### Anti-KDM5C / Jarid1C / SMCX antibody ab34718

KO 評価済

★★★★☆ 5 Abreviews 12 References 画像数 6

#### 製品の概要



製品名	Anti-KDM5C / Jarid1C / SMCX antibody
製品の詳細	Rabbit polyclonal to KDM5C / Jarid1C / SMCX
由来種	Rabbit
アプリケーション	適用あり: WB, ICC/IF
種交差性	交差種: Human 交差が予測される動物種: Pig 
免疫原	Synthetic peptide conjugated to KLH derived from within residues 1500 to the C-terminus of Human Jarid1C/ SMCX.Immunogen の所有権に関して(Peptide available as <a href="#">ab35501</a> .)
ポジティブ・コントロール	WB: HEK-293T, HAP1 and Y79 cell lysates. ICC/IF: HeLa cells.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

#### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	pH: 7.40 Preservative: 0.02% Sodium azide Constituent: PBS  Batches of this product that have a concentration < 1mg/ml may have BSA added as a stabilising agent. If you would like information about the formulation of a specific lot, please contact our scientific support team who will be happy to help.
精製度	Immunogen affinity purified


ポリクローナル  
IgG

**The Abpromise guarantee**      Abpromise保証は、次のテスト済みアプリケーションにおけるab34718の使用に適用されます  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB	 (3)	1/250. Detects a band of approximately 180 kDa (predicted molecular weight: 176 kDa). Abcam recommends using milk as the blocking agent.
ICC/IF	 (1)	Use a concentration of 1 µg/ml.

機能	Histone demethylase that specifically demethylates 'Lys-4' of histone H3, thereby playing a central role in histone code. Does not demethylate histone H3 'Lys-9', H3 'Lys-27', H3 'Lys-36', H3 'Lys-79' or H4 'Lys-20'. Demethylates trimethylated and dimethylated but not monomethylated H3 'Lys-4'. Participates in transcriptional repression of neuronal genes by recruiting histone deacetylases and REST at neuron-restrictive silencer elements.
組織特異性	Expressed in all tissues examined. Highest levels found in brain and skeletal muscle.
関連疾患	Defects in KDM5C are the cause of mental retardation syndromic X-linked JARID1C-related (MRXSJ) [MIM:300534]. MRXSJ is characterized by significantly sub-average general intellectual functioning associated with impairments in adaptive behavior and manifested during the developmental period. MRXSJ patients manifest mental retardation associated with variable features such as slowly progressive spastic paraplegia, seizures, facial dysmorphism.
配列類似性	Belongs to the JARID1 histone demethylase family. Contains 1 ARID domain. Contains 1 JmjC domain. Contains 1 JmjN domain. Contains 2 PHD-type zinc fingers.
ドメイン	The first PHD-type zinc finger domain recognizes and binds H3-K9Me3. Both the JmjC domain and the JmjN domain are required for enzymatic activity.
細胞内局在	Nucleus.

## 2



**Lane 1:** Wild-type HAP1 whole cell lysate (20 µg)


**Lane 2:** KDM5C knockout HAP1 whole cell lysate (20 µg)

**Lane 3:** HEK293 whole cell lysate (20 µg)

**Lane 4:** U2OS whole cell lysate (20 µg)


**Lanes 1 - 4:** Merged signal (red and green). Green - ab34718 observed at 175 kDa. Red - loading control, **ab18058**, observed at 120 kDa.

ab34718 was shown to specifically recognize KDM5C in wild-type HAP1 cells as signal was lost at the expected MW in KDM5C knockout cells. Additional cross-reactive bands were observed in the wild-type and knockout cells. Wild-type and KDM5C knockout samples were subjected to SDS-PAGE. Ab34718 and **ab18058** (Mouse anti Vinculin loading control) were incubated overnight at 4°C at 1/250 dilution and 1/20,000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed (**ab216776**) secondary antibodies at 1/20,000 dilution for 1 hour at room temperature before imaging.




ab34718 staining KDM5C / Jarid1C / SMCX in HeLa cells. The cells were fixed with 100% methanol (5 min), permeabilized with 0.1% PBS-Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at 4°C with ab34718 at 1µg/ml and **ab7291**, Mouse monoclonal [DM1A] to alpha Tubulin - Loading Control. Cells were then incubated with **ab150081**, Goat polyclonal Secondary Antibody to Rabbit IgG - H&L (Alexa Fluor® 488), pre-adsorbed at 1/1000 dilution (shown in green) and **ab150120**, Goat polyclonal Secondary Antibody to Mouse IgG - H&L (Alexa Fluor® 594), pre-adsorbed at 1/1000 dilution (shown in pseudocolour red). Nuclear DNA was labelled with DAPI (shown in blue).

Image was acquired with a high-content analyser (Operetta CLS, Perkin Elmer) and a maximum intensity projection of confocal sections is shown.



Immunocytochemistry/ Immunofluorescence - Anti-KDM5C / Jarid1C / SMCX antibody (ab34718)



Western blot - Anti-KDM5C / Jarid1C / SMCX antibody (ab34718)

PFA-fixed, 0.5% Triton X-100 permeabilized HeLa (human epithelial cell line from cervix adenocarcinoma) cells stained for KDM5C / Jarid1C / SMCX (green) using ab34718 at 1/200 dilution in ICC/IF. Counter-stained with DAPI in order to highlight the nucleus (red). Please refer to abreview for further experimental details.

**All lanes :** Anti-KDM5C / Jarid1C / SMCX antibody (ab34718) at 1/250 dilution

**Lane 1 :** Wild-type HEK-293T cell lysate

**Lane 2 :** KDM5C knockout HEK-293T cell lysate

Lysates/proteins at 20 µg per lane.


Performed under reducing conditions.

**Predicted band size:** 176 kDa

**Observed band size:** 175 kDa

**Lanes 1 -2:** Merged signal (red and green). Green - ab34718 observed at 175 kDa. Red - loading control **ab7291** (Mouse anti-Alpha Tubulin [DM1A]) observed at 55kDa.

ab34718 was shown to react with KDM5C / Jarid1C / SMCX in wild-type HEK-293T cells in western blot with loss of signal observed in KDM5C knockout cell line **ab266251** (KDM5C knockout cell lysate **ab257494**). Wild-type and KDM5C knockout HEK-293T cell lysates were subjected to SDS-PAGE. Membranes were blocked in 3% milk in TBS-T (0.1% Tween®) before incubation with ab34718 and **ab7291** (Mouse anti-Alpha Tubulin [DM1A]) overnight at 4°C at a 1 in 250 dilution and a 1 in 20000 dilution respectively. Blots were incubated with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed (**ab216776**) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



Western blot - Anti-KDM5C / Jarid1C / SMCX antibody (ab34718)

**All lanes :** Anti-KDM5C / Jarid1C / SMCX antibody (ab34718) at 1 µg/ml

**Lane 1 :** HEK293 (Human) Whole Cell Lysate

**Lane 2 :** Y79 (Human retinoblastoma cell line) Whole Cell Lysate

Lysates/proteins at 20 µg per lane.

### Secondary

**All lanes :** Goat Anti-Rabbit IgG H&L (HRP) preadsorbed at 1/50000 dilution

Developed using the ECL technique.

Performed under reducing conditions.


**Predicted band size:** 176 kDa

**Observed band size:** 180 kDa

**Additional bands at:** 45 kDa, 58 kDa. We are unsure as to the identity of these extra bands.

**Exposure time:** 20 minutes

This blot was produced using a 3-8% Tris Acetate gel under the TA buffer system. The gel was run at 150V for 60 minutes before being transferred onto a Nitrocellulose membrane at 30V for 70 minutes. The membrane was then blocked for an hour using 3% Milk before being incubated with ab34718 overnight at 4°C. Antibody binding was detected using an anti-rabbit antibody conjugated to HRP, and visualised using ECL development solution [\*\*ab133406\*\*](#).



Anti-KDM5C / Jarid1C / SMCX antibody (ab34718) at 1/250 dilution + HEK-293 whole cell lysate (**ab7902**) at 20 µg

#### Secondary

IRDye 680 Conjugated Goat Anti-Rabbit IgG (H+L) at 1/10000 dilution

Performed under reducing conditions.

**Predicted band size:** 176 kDa

**Observed band size:** 176 kDa

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

#### Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors